



**Europäisches  
Patentamt**

**European  
Patent Office**

**Office européen  
des brevets**

*EP03/14679*

REC'D 06 FEB 2004	
WIPO	PCT

**Bescheinigung**

**Certificate**

**Attestation**

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

**Patentanmeldung Nr. Patent application No. Demande de brevet n°**

03004394.7

BEST AVAILABLE COPY

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

**R C van Dijk**



Anmeldung Nr:  
Application no.: 03004394.7 ✓  
Demande no:

Anmeldetag:  
Date of filing: 03.03.03 ✓  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Julius-Maximilians-Universität Würzburg  
Sanderring 2  
97070 Würzburg  
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Millisecond activation switch for seven-transmembrane proteins

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)  
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt: ..

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

C07K14/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PT SE SI SK TR LI

EPO • Munich  
67  
03. März 2003

New EP patent application  
Universität Würzburg  
Our Ref.: H 1340 EP

### **Millisecond activation switch for seven-transmembrane proteins**

The present invention relates to recombinant seven-transmembrane receptor, whereby the amino terminus of said recombinant receptor is located on an extracellular side and the carboxy-terminus is located on an intracellular side of a membrane, comprising at least two detectable labels, whereby a first of said at least two detectable labels is or is located (on) the carboxy-terminus and whereby a second of said at least two labels is or is located (on) the first or third intracellular loop or whereby a first of said at least two labels is or is located (on) the first intracellular loop and a second of at said at least two labels is or is located (on) the third intracellular loop. Furthermore, nucleic acid molecules encoding said recombinant seven-transmembrane receptors are described as well as vector and host cells comprising the same. Furthermore, the present invention provides for identification and screening methods for molecules or compounds which are capable of modifying the biological end of pharmacological function of seven-transmembrane receptor proteins. Finally, diagnostic compositions comprising the compounds of the present invention as well as kits comprising said compounds are disclosed.

Several seven-transmembrane proteins have been described within the last twenty years, whereby such proteins are mainly involved in signal transduction pathways. Most of these seven-transmembrane proteins are to be classified as proto-oncogenes or as specific receptors which are commonly linked to a signal transduction pathways involving G proteins. G protein-coupled seven transmembrane segment receptors (G protein-coupled receptors, GPCRs, or 7TM receptors) are a very heterogenous class of molecules comprising more than 1000 different members. Accordingly, GPCRs are one of the largest superfamilies

of proteins in the human and mammalian body. GPCRs do not share any overall sequence homology yet, one characteristic structural feature, common to all GPCRs is the presence of the characteristic seven-transmembrane spanning  $\alpha$ -helical segment connected by alternating intracellular and extracellular loops, whereby the amino terminal is located on the extracellular side and the carboxy terminus is located on the intracellular side. As summarized in Gether, *Endocrine Rev.* 21 (2000), 90 to 113, G protein-coupled receptors can be classified in several families, termed family "A to C" or family "1 to 3". Family A is normally classified as the rhodopsin/ $\beta$ 2-adrenergic receptor-like GPCRs. This family again is subdivided phylogenetically into six subgroups, namely 1. biogenic amine receptors (adrenergic serotonin, dopamine, muscarinic, histamine); 2. CCK-, endothelin-, tachykinin-, neuropeptide Y-, TRH-, neurotensin-, bombesin- and growth hormone secretagogues-receptors plus vertebrate opsin(s)-receptors; 3. invertebrate opsins- and bradykinin-receptors; 4. adenosine-, cannabinoid-, melanocortin- and biractory-receptors; 5. chemokine-, fMLP-, C5A-, GnRH-, eicosanoid-, leukotriene-, FSH-, LH-, TSH-, fMLP-, galanin-, nucleotide opioid-, oxytocin-receptors; as well as 6. melatonin receptors and other non-classified receptors. Family B is the family related to glucagons/VIP/calcitonin receptor-like GPCRs which comprise four GPCR-subfamilies, namely: 1. calcitonin-, CGRP- and -receptors; 2. PTH- and PTHrP-receptors; 3. glucagons-, glucagon-like peptide-, GIP-, GHRH-, PACAP-, VIP- and secretin-receptors as well as 4. latrotoxin-receptor. Finally, family C is characterized as the metabotropic neurotransmitter/calcium GPCRs. In this family the following receptors are, inter alia, comprised: 1. metabotropic glutamate-receptors; 2. metabotropic GABA-receptors; 3. calcium-receptors; 4. vomeronasal pheromone receptors; and 5. taste receptors.

As mentioned above, G-protein-coupled receptors (GPCRs) are the largest family of hormone or neurotransmitter receptors; they have a common structure containing seven transmembrane  $\alpha$ -helices (Rohrer, *Physiol. Rev.* 78 (1998), 35-52; Gether, *Endocr. Rev.* 21 (2000) 90-113; Pierce, *Mol. Cell Biol.* 3 (2002), 639-650). Their activation by specific agonists – hormones or neurotransmitters –



switches them into an active state that couples to and activates G-proteins, the signal transducers. The G-proteins, in turn, can activate a multitude of effector proteins such as ion channels or 2<sup>nd</sup> messenger producing enzymes that alter many functions (e.g., cardiovascular, neural, endocrine) in virtually every type of cells. A large body of data suggests that agonist-induced activation leads to a relative rearrangement of the receptor's transmembrane helices, most notably of helix III and VI (Farrens, *Science* 274 (1996), 768-770; Sheikh, *Nature* 383 (1996), 347-350; Sheikh, *J. Biol. Chem.* 274 (1999), 17033-17041).

A very special GPCR is the "light receptor" rhodopsin that senses light by using a covalently coupled ligand, 11-cis retinal, that isomerizes upon capture of a photon. In this case, conformational changes in the "receptor" protein can be inferred from spectroscopic studies of the bound retinal, and multiple activation states formed within milliseconds have been described (Farrens, (1996), loc. cit.); Okada, *Trends Biochem. Sci.* 26 (2001), 318-324). No comparable techniques are available for hormone- or transmitter-activated receptors. Spectroscopic studies have been done with purified  $\beta_2$ -adrenergic receptor chemically labelled with fluorophores and reconstituted into lipid membranes (Gether, *J. Biol. Chem.* 270 (1995), 28268-28275; Jensen, *J. Biol. Chem.* 276 (2001), 9279-9290; Ghanouni, *J. Biol. Chem.* 276 (2001), 24433-24436; Ghanouni, *Proc. Natl. Acad. Sci. USA* 98 (2001), 5997-6002). These studies observed agonist-mediated fluorescence changes in the minute time scale. This is much slower than biological responses to receptor activation, which can occur within seconds.

The advent of fluorescent proteins has allowed non-invasive intracellular labeling, especially of peptides, which are easily detectable by optical means. The green fluorescent protein (GFP) from *Aequorea victoria* is now the most widely used reporter gene in many organisms. Multiple variants with different spectral properties have been developed. Furthermore, combinations of fluorescent proteins exhibiting energy transfer provide for differential fluorescence in response to conformational changes in the protein's immediate environment. Based on this principle, fusion constructs have been developed which allow to detect specific analytes such as calcium ions, cAMP or cGMP (e.g. WO 98/40477; Honda, *Proc.*

Natl. Acad. Sci. USA 98 (2001), 2437-2442; Zaccolo, Nat. Cell. Biol. 2 (2000), 25-29). Furthermore, in recent years, detection methods for monitoring fluorescence resonance energy transfer (FRET) have been described for verifying and/or detecting homo- or heterodimerization of molecules or receptor-mediated activation processes. US 02/0048811 describes the visualization of receptor-mediated activation of heterotrimeric G-proteins by FRET. Here, the association and/or activation of pairs of G-proteins or of corresponding subunits were measured. DE-A1 101 08 263 relates to the analysis of modifications of proteins with ubiquitin-related proteins. DE-A1 101 08 263 provides a FRET-based test-system comprising ubiquitin-related donor parts and acceptor parts which relates to corresponding target molecules. Both patent applications, US 02/0048811 and DE-A1 101 08 263 exemplify the large body of literature relating to FRET-analysis of "two-component-assays". These assays relate to the distinct use of independent "donor" and "acceptor" molecules/moieties.

Some further limited approaches have been undertaken to employ FRET-technology on an intramolecular basis. For instance, WO 98/40477 describes such a system based on calmodulin which is useful for the measurement of calcium concentrations. However, this system requires as a second analyte binding portion a calcium-calmodulin binding target peptide moiety to which the conformationally flexible calmodulin binds. Thus, the applicability of the fusion protein of WO 98/40477 is restricted because of the requirement of two binding portions and the use of a ubiquitous and endogenous regulatory component.

US 6,277,627 relates to a glucose biosensor comprising at least one fluorophore and in US 6,197,534 a glucose biosensor is provided which comprises a glucose binding protein with two fluorophores to be employed in FRET-like technologies. Yet, these documents provide for fluorophore-labeled proteins of rather small size which are soluble.

However, the above discussed seven-transmembrane proteins, in particular proto-oncogenes or G protein-coupled receptors (GPCRs) are rather large ("bulky")

membrane proteins, wherein FRET- or BRET-based assays have merely been described in dimerization studies.

Based on the large number of GPCR family members and the prominent role in regulating cellular signals, GPCRs represent the most important family of drug targets. The following methods are available to study interaction of potential ligands with GPCRs: a. Ligand-binding assay: This method is restricted to available radio- or fluorescent-labelled ligands, which limits its use to known receptors. Based on the nature of this assay, just the interaction of the ligand with the receptor can be studied but no information regarding the activating, blocking- or inhibiting properties of the ligand on the receptor can be gathered. b. Recording of the activity of G-protein effector systems has become the most important method for drug screening of GPCRs. (Milligan G & Rees S; Trends Pharmacol Sci. 1999 Mar;20(3):118-24.) To measure the activity of G-protein effectors makes high-throughput screening on cell based assays possible, however because of the fact that the effector system is several steps downstream of the receptor activation this method has following disadvantages: 1. It is prone to unspecific drug effects, that are not mediated via the investigated GPCR, but rather result from either interaction with elements of the signalling cascade that are downstream of GPCRs or are mediated in parallel via other (endogenous) GPCRs of which many kind are present in various cell systems ( <http://www.tumor-gene.org/GPCR/gpcr.html>.) 2. Receptor activation and deactivation cannot be determined in real time. Therefore, it is impossible to distinguish between receptor activation and fast receptor desensitization. 3. Recording of GPCR activity depends on expression levels and specificity of subsequent G-proteins and effectors, preventing in many cases exact comparisons between different GPCR subtypes. 4. The strength of the signal or the potency of a ligand to induce full activation of a cellular signal will largely depend on the expression level of the receptor (Bünemann et al. J Biol Chem. 2001 Dec 14;276(50):47512-7.). Uncontrolled fluctuations of the expression level will cause variability of the result, and will again make comparisons between different GPCR subtypes difficult. 5. A possibility to

measure agonism, partial agonism, inverse agonism and neutral antagonism on the level of the receptor is lacking.

Due to these disadvantages, novel methods that allow detection of receptor activity at the level of the receptor have been tried: Subsequent to the discovery that many GPCRs dimerize or oligomerize it was tried to detect agonist-mediated conformational changes of GPCRs via detection of changes in inter-molecular BRET or FRET (Angers, *Annu Rev Pharmacol Toxicol* 2002;42:409-35). However, in the vast majority of tested receptors no ligand-induced change in BRET or FRET could be detected (Angers, (2002), loc. cit.; Rios, *Pharmacol Ther* 2001 Nov-Dec;92(2-3):71-87).

A second potential method to monitor GPCR activation based on fluorescence-quenching was developed by using  $\beta$ -2 adrenergic receptors which were mutated by eliminating all but one accessible cysteines and after purification were labelled with a fluorescent tag. In reconstituted membranes, this method was the first to detect ligand induced conformational changes in a GPCR in real time, however the obtained activation kinetics of these receptors were too slow to be in accordance with physiological responses induced via  $\beta$ 2-adrenergic receptors (Kobilka & Gether, *Methods Enzymol.* 2002;343:170-82).

As mentioned above the GPRC family as well as seven-transmembrane proto-oncogene provide for highly interesting target in drug development. However, the prior art as discussed above provided merely for test system, wherein antagonists and agonists could be measured in unphysiological time frames. In particular, studies as described above observe agonist mediated changes on, for example, purified  $\beta$ 2-adrenergic receptors in the minute time scale.

Therefore, apart from the requirements for detecting specific modifiers for seven-transmembrane proteins, in particular proto-oncogenes or GPCRs, there is also a need for means and methods that allow reliable, fast and easy measurement of the activation of such seven-transmembrane proteins. Such measurements have

not been provided for or are not yet accessible by prior art techniques. Preferably, the desired means and methods allow for in vivo measurements.

This technical problem is solved by the provisions of the embodiments as characterized in the claims.

Accordingly, the present invention relates to a recombinant seven-transmembrane receptor, whereby the amino terminus of said recombinant receptor is located on an extracellular side and the carboxy-terminus is located on an intracellular side of a membrane, comprising at least two detectable labels, whereby a first of said at least two detectable labels is or is located (on) the carboxy-terminus and whereby a second of said at least two labels is or is located (on) the first or third intracellular loop or whereby a first of said at least two labels is or is located (on) the first intracellular loop and a second of at said at least two labels is or is located (on) the third intracellular loop.

Inter alia, hormones and neurotransmitters transduce signals via, seven-transmembrane proteins G-protein-coupled receptors as key switches in order to change cellular functions. Despite utilizing common signalling pathways, hormone, amino acids, drugs and neurotransmitter responses exhibit different temporal patterns. To reveal the molecular basis for such differences, the present invention provides for a generally applicable fluorescence-based technique for real-time monitoring of the activation switch for G protein-coupled receptors in single cells, liposomes or membrane preparations. Such direct measurements as disclosed herein were used to investigate the activation of three exemplified seven-transmembrane proteins, namely the  $\alpha_{2A}$ -adrenergic (neurotransmitter) receptor, the (adenosine) A<sub>2A</sub>-receptor and the parathyroid hormone (PTH, hormone) receptors. Surprisingly, kinetics that were much faster than previously thought could be observed, for example,  $\approx 40$  ms for the  $\alpha_{2A}$ -adrenergic receptor and  $\approx 1$  s for the PTH receptor. The different switch times are in agreement with the distinct biological functions of these receptors. Agonist, partial agonist and antagonist may rapidly switch on and off the receptors in proportion to their

respective intrinsic activities. The means and methods provided herein permit the comparison even of agonists and partial agonists of intrinsic activities at the receptors or proto-oncogenes themselves and allow for the detection of millisecond activation-times of G-protein coupled receptors (GPCRs) as well as proto-oncogenes.

Previous work has shown that seven-transmembrane proteins in particular GPCR comprise an entire surface of the cytoplasmic part of about 40 Ångstrom (Bourne, Science 289 (2000), 733-4 and Teller, Biochemistry 40 (2001), 7768-7772). Yet, fluorophores, like GFP variants show a cylindrical or rod-like structure with a diameter of 30 Ångstrom. Accordingly, it was expected that insertion of two fluorophores into the intracellular parts of a seven-transmembrane protein, in particular a GPCR, would cover the entire surface of the cytoplasmic part of said protein or receptor and thereby prevent interaction with protein-coupled further signal transduction part of receptors like arrestins, kinases, or gene proteins. Here, it was surprisingly found that the use of two GFP analogs or fluorophores did not constrain or alter the movements of the transmembrane helices and the receptor was still functional and even provides for a test system wherein intramolecular movements of seven-transmembrane proteins can be monitored in a millisecond range. Therefore, it was surprisingly found that despite the fact that the size of even one GFP variant (diameter of 30 Ångstrom, (Tsien, Annu. Rev. Biochem. 1998. 67:509-544) covers almost the entire intracellular surface of a GPCR, receptor constructs fused to two GFP variants are still capable to activate heterotrimeric G proteins.

This is in particular surprising, since it was envisaged that fusing large protein moieties, like YFP or GFP, to GPCRs constrains or grossly alters movements of transmembran helices of the receptor required for receptor activation.

The prior art has provided for modifications of GPCRs, for example for point mutation in the 3<sup>rd</sup> intracellular loop located after the 5<sup>th</sup> transmembrane region; see Huang, JBC 271 (1996), 33382-33389. However, in this study the introduced mutation caused dramatic changes in the activity of the GPCR studied. Similarly,

modifications of proteins involved in signal transduction by incorporation of, inter alia, fluorophores often leads to complete inactivation of the additional protein. As documented in the appended examples, it could be shown that the introduction of green fluorescence protein on position Pro116 of the  $\alpha A$ - $\alpha B$  loop of G protein receptor-interacting  $G\alpha$  completely inactivated said interacting protein and correspondingly, the signal transduction pathway. Similarly, Belke-Louis and Schulz (Naunyn-Schmiedeberg's Arch. Pharmacol. 2000, 391(suppl.), R51 (189) reported N-terminal or C-terminal fusion proteins of  $G\alpha s$  with EGFP exhibited dramatically impaired functionality.

Furthermore, the expected actual movement of transmembrane helices in GPCRs upon activation by agonists are very small, in particular in the range of 2 to 8 Ångström (Altenbach, Biochemistry 25 (2001), 15493-15500). Accordingly, it could certainly not be expected that the recombinant membrane receptors as defined herein provide for assay systems which are capable of detecting this minor conformational changes and movements by FRET- or BRET-technology, employing fluorophores or chromophores which are located at distal of the agonist activation side.

Accordingly, it could not be expected that the preparation of a recombinant seven-transmembrane receptor as described herein provides for a fast-kinetic, functional and reliable tool for the measurements of receptor-activation or -inhibition.

The term "recombinant seven-transmembrane receptor" as defined herein relates to a membrane protein which is recombinantly produced by methods known in the art; see, inter alia, Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2001. The "recombinant seven-transmembrane receptor" is preferably inserted into a biological and/or artificial membrane, like a cellular membrane, a crude membrane preparation, liposomes as well as artificial membranes comprising micelles, lipid monolayer or lipid bilayers. Yet, most preferably, in accordance with this invention, the recombinant seven-transmembrane receptors as defined herein are located in cellular membranes, for example membranes of cultured cells or membranes of

oocytes. As detailed herein below, it is also envisaged that the recombinant seven-transmembrane receptors as defined herein are expressed in transgenic, non-human animals. Accordingly, also cells, tissues and organs of said non-human transgenic animal may express the recombinant seven-transmembrane receptors of the present invention and may be in particular useful in drug screenings.

The term "recombinant seven-transmembrane receptor" also relates to a seven-transmembrane protein, recombinantly produced and comprising said at least two detectable labels, whereby said recombinantly produced membrane receptor/membrane protein may also consist of a chimeric receptor molecule, i.e. a receptor molecule which is derived from two or more naturally occurring seven-transmembrane proteins. It is, inter alia, envisaged that seven-transmembrane receptors are engineered and recombinantly produced which comprise parts of one GPCR and further parts of another GPCR.

The term "and/or" wherever used herein includes the meaning of "and", "or" all or any other combination of the elements connected by said term.

The terms "amino terminus" or "carboxy terminus" of the recombinant seven-transmembrane receptor corresponds to the amino terminus and carboxy terminus of non-modified naturally occurring seven-transmembrane receptors or proto-oncogenes. The corresponding topology and its verification is, inter alia, disclosed in Gether (2000), loc. cit. Accordingly, the natural orientation of seven-transmembrane receptors is known in the art as, for example, shown in Gether (2000), loc. cit. The orientation also relates to the topology of the inventive recombinant receptors in (crude) membrane preparations or liposomes. Accordingly, the N-terminus should in liposomes be oriented to the outside, whereas the C-terminus should be in the lumen of the liposomes, membrane vesicles. Furthermore, the person skilled in the art is aware of suitable techniques for detecting the orientation of a transmembrane protein and, accordingly, of the recombinant seven-transmembrane receptor of the present invention. Such



techniques comprise but are not limited to crystallography, NMR-studies, modeling studies as well as microscopy techniques, like immunolabeling in electron microscopy preparation and the like.

The term "membrane" as used herein and in particular in context of the recombinant seven-transmembrane receptors/proteins of the present invention relate to naturally occurring membranes as well as to artificial membranes. Preferably, the membranes consist of lipid bilayers. As pointed out above, specific examples are cellular membranes and bio-membranes, like the plasma membrane of cells, the endoplasmic reticulum, mitochondrial membrane, golgi vesicles, lysosomes, peroxisomes, but also cellular membranes of plant cells, like membranes of the chloroplasts or other organelles as well as vacuoles. Yet, most preferably, the cellular or bio-membrane into which the recombinant protein/receptor of the invention is inserted is the plasma membrane of an animal cell, most preferably of a mammalian cell, but also of amphibian cells, like frog oocytes. Yet, as also discussed herein, membrane preparations, like crude membrane preparations or liposomes are envisaged as "membranes" wherein the recombinant membrane receptor/protein of the present invention is inserted.

The term "at least two detectable labels" as used herein means that the recombinant membrane receptor/protein of the invention may comprise at least two, but also three or more detectable labels. The detectable labels will be detailed herein below and may, in particular comprise fluorophores as well as bioluminescent substances. In accordance with the appended examples, however, most preferred are two detectable labels on one recombinant seven-transmembrane protein of the invention.

The term "said label is or is located (on)" as employed herein means that the label may either be part of the corresponding region of the recombinant seven-transmembrane receptor, i.e. the first or third intracellular loop or of the carboxy terminus or said label may completely replace the corresponding region of the recombinant seven-transmembrane receptor/protein, i.e. the first or the third

intracellular loop may be replaced by the corresponding label or the C-terminus may be replaced by the corresponding label. Yet, as documented in the appended examples and herein below, it is preferred that the labels are introduced in a region comprised in the first or the third intracellular loop or said at least one label replaces only parts of the C (carboxy)-terminus of the recombinant seven-transmembrane receptor. Accordingly, it is preferred that at least one of said labels is introduced within the first or third intracellular loop of the naturally occurring seven-transmembrane receptor, preferably in naturally occurring GPCRs. Some corresponding third intracellular loop regions are exemplified in the examples and comprise, inter alia, the amino acid sequences as shown in SEQ ID NO: 18, 22 or 26. These third intracellular loop regions may be encoded by sequences as depicted in SEQ ID NOS: 17, 21 and 25. Corresponding exemplified C-terminal region which may comprise one of the detectable labels are shown in form of amino acid sequences in SEQ ID NOS: 19, 23 and 28. In particular, SEQ ID NO: 28 depicts the carboxy terminus of human PTH/PTHrP receptor corresponding modifications of this C-terminus are exemplified in SEQ ID NOS: 29 to 37. Here, in this context, parts of the C-terminus have been deleted and replaced with a green fluorescent protein variant, namely YFP (yellow fluorescent protein). Preferred sites for introduction of the detectable labels are illustrated below and in the appended examples. Furthermore, the examples provide for naturally occurring seven-transmembrane receptors like the  $\alpha$ 2A-adrenergic receptor, the PTH/PTHrP receptor or the A2A-adenosine receptor which are modified by recombinant technology and comprise at least two detectable labels which are, in particular located on (or within) the third intracellular loop. Corresponding examples are shown in SEQ ID NOS: 12, 14 and 16. In a preferred embodiment of the present invention the detectable label is inserted into the third intracellular loop of the seven-transmembrane receptor/protein and preferably after the second amino acid of the transmembrane region 5 (TM5) and before the second amino acid before the beginning of transmembrane region 6 (TM6). Even more preferred, the detectable label is inserted after amino acid 8 after transmembrane region 5 (TM5) or after amino acid 22 after transmembrane region 5 (TM5). In a preferred embodiment the

detectable label is inserted 10 or 12 amino acids before the transmembrane region 6 (TM6). As far as the insertion of the detectable label on the carboxy terminus is concerned, it is preferred that 5 to 25 amino acids of the natural carboxy terminus remain at the end of transmembrane region 7 (TM7). Preferably, the detectable label is inserted after the 16<sup>th</sup> or 20<sup>th</sup> amino acid after transmembrane region 7 (TM7). Therefore, it is envisaged that the detectable labels either replace naturally occurring amino acids on the first and/or third loop of seven-transmembrane receptors or the C-terminus of the same or that said detectable labels are inserted into the naturally occurring amino acid sequences of the same.

As far as insertion of detectable labels of the first intracellular loop are concerned, again, it is preferred that said label is inserted two amino acids after the end of transmembrane region 1 (TM1) and two amino acids before the beginning of transmembrane region 2 (TM2). Most preferably, the detectable label is inserted in the middle of the first intracellular loop. The same applies, *mutatis mutandis* for the third intracellular loop insertion. Yet, it is to be noted that all insertions of the detectable label may lead to a deletion/replacement of naturally occurring amino acids in the first or third intracellular loop as well as the C-terminus.

In a most preferred embodiment of the recombinant membrane receptor of the invention, the first detectable label is or is located (on) the third intracellular loop of said membrane receptor and the second detectable label is or is located (on) the carboxy terminus. This most preferred embodiment of the recombinant membrane receptor/protein of the present invention is also exemplified in the appended scientific examples. Furthermore, corresponding recombinant constructs are depicted as nucleic acid sequences in appended SEQ ID NOS: 11, 13 and 15 and in the corresponding amino acid sequences SEQ ID NOS: 12, 14 and 16. However, it is of note that these amino acid sequences as well as the encoding nucleic acid sequences merely comprise illustrative examples.

The invention in particular relates to a recombinant membrane receptor/protein, whereby said membrane receptor is a G-protein-coupled receptor or a proto-oncogene. Accordingly, the present invention provides for recombinant membrane receptors which are derived from naturally occurring G protein-coupled receptors or naturally occurring proto-oncogenes. As shown in the appended examples the invention provides for genetically/recombinantly modified G protein-coupled receptors or proto-oncogenes which are particularly useful in methods provided herein, i.e. screening methods for antagonists as well as agonists of G protein-coupled receptors or proto-oncogenes, preferably involved in the signal transduction pathways. The proto-oncogenes which are in their naturally occurring form seven-transmembrane proteins and which are, in accordance with this invention, modified and recombinantly produced comprise the above-recited at least two detectable labels on the above-recited positions. In a preferred embodiment this proto-oncogene belongs to the "frizzled/smoothened family" and comprises the protein "frizzled" or "smoothened".

In a most preferred embodiment of the present invention the recombinant seven-transmembrane receptor is derived from a naturally occurring G protein-coupled receptor or is derived from a corresponding orphan receptor; i.e. putative or unclassified GPCRs. G protein-coupled receptors to be modified and recombinantly produced in accordance with this invention comprise the GPCRs as, inter alia, defined in Gether (2000), loc. cit. as well as GPCRs comprised in the following classes A to E. Further general classifications of the GPCR superfamily are provided by the classification in class A to E.

Class A (or class 1) comprises rhodopsin-like receptors family, class B (or class 2) comprises secretin-like receptors family, class C (or class 3) comprises calcium sensing-like receptors family, class D (or class 4) comprises yeast alpha factor-like receptors family, class E (or class 5) comprises dictyostelium attractant-like receptors family.

Class A rhodopsin like GPCRs comprise, but are not limited to amine-, peptide-, hormone protein-, (rhod)opsin-, olfactory-, prostanoid-, nucleotide-like-, cannabis-, platelet activating factor-, gonadotropin-releasing hormone-, thyrotropin-releasing hormone & secretagogue-, melatonin-, viral-, lysosphingolipid & LPA (EDG)-, leukotriene B4 receptor, class A Orphan/other receptor.

Class B Secretin like GPCRs comprise, but are not limited to calcitonin-, corticotropin releasing factor-, gastric inhibitory peptide-, glucagon-, growth hormone-releasing hormone-, parathyroid hormone-, PACAP-, secretin-, vasoactive intestinal polypeptide-, diuretic hormone-, EMR1-, latrophilin-, brain-specific angiogenesis inhibitor (BAI)-, methuselah-like proteins (MTH)-, cadherin EGF LAG (CELSR)-receptor.

Class C Metabotropic glutamate / pheromone like GPCRs comprise, but are not limited to metabotropic glutamate-, extracellular calcium-sensing-, putative pheromone receptors, GABA-B-, orphan GPRC5-receptor.

Class D Fungal pheromone comprise, but are not limited to fungal pheromone STE2-like-, fungal pheromone STE3-like-receptor.

Furthermore, the invention also provides for modified GPCRs from class E, namely cAMP receptors (Dictyostelium), like frizzled/Smoothed family-, frizzled-, Smoothed-receptor.

Accordingly, the present invention is not limited to a modified or recombinantly produced seven-transmembrane receptors of mammalian origin but also comprise recombinantly seven-transmembrane receptors from other eukaryotes, including yeasts, plants, fungi as well as non-vertebrates animals.

The recombinant membrane receptor/protein of the invention is, preferably derived from a naturally occurring proto-oncogene like smoothed receptor (Smo) or frizzled receptor or is a G-protein-coupled receptor (GPCR) is selected

from the group consisting of a rhodopsin/ $\beta$ 2 adrenergic receptor-like GPCR, a glucagon/VIP/calcitonin receptor-like GPCR and a metabotropic neurotransmitter/calcium receptor. In a most preferred embodiment, said rhodopsin/ $\beta$ 2-adrenergic receptor-like GPCR is the  $\alpha$ 2A adrenergic receptor or the adenosine receptor A2A and said glucagon/VIP/calcitonin receptor-like GPCR is the parathyroid hormone (PTH) receptor.

In accordance with this invention and as pointed out herein above, naturally occurring seven-transmembrane protein may be genetically/recombinantly modified in order to provide from the recombinant seven-transmembrane receptors of the present invention. For example, the human PTH/PTHrP receptor gene is well known in the art and deposited under GeneBank accession No. U22401. A mouse  $\alpha$ 2A-adrenergic receptor can be found under GeneBank accession No. M99377. The gene encoding for human A2A receptor can be found in GeneBank accession No. M97370. Corresponding sequences on nucleic acid as well as amino acid level are depicted and described in appended SEQ ID NOS: 1 to 6. Yet, it is of note that the G protein-coupled receptors or the proto-oncogene to be employed in accordance with the present invention for the generation of recombinant seven-transmembrane receptors as defined herein above are not limited to GPCRs or proto-oncogenes derived from human or mouse. It is in particular settings also envisaged to employ G protein-coupled receptors or proto-oncogenes from other vertebrates like rats, rabbits, guinea pigs, dogs, cats or frogs and fish.

The detectable labels to be introduced into the recombinant seven-transmembrane receptor/proteins of the present invention are preferably fluorescent labels or bioluminescent labels.

The fluorescence labels are, in particular, selected from "green fluorescent protein" or "GFP" and its variants or mutants. Preferably, said group comprises GFP (green), YFP (yellow), CFP (cyan), BFP (blue) and dsRed.

The bioluminescent labels may be luciferase (like renilla luciferase or firefly luciferase). Furthermore, it is envisaged that the fluorescence label is produced by binding the Flash compound to specific epitopes of said 1<sup>st</sup> and 3<sup>rd</sup> loop or said C-terminus of the recombinant seven-transmembrane receptor.

FLASH (fluorescein arsenical helix binder) can specifically bind a protein comprising the sequence Cys-Cys-X-X-Cys-Cys (X represents any amino acid but preferentially X-X is Pro-Gly). FLASH fluoresces after binding to the sequence Cys-Cys-X-X-Cys-Cys, and thus allows site-specific fluorescent labeling of recombinant proteins in living cells. Accordingly, in context of the present invention, intramolecular FRET with GPCRs may be performed with a recombinant receptor containing the bound FLASH (to the epitope, Cys-Cys-X-X-Cys-Cys ) and the, inter alia, CFP; see, Griffin, Science 281 (1998), 269-272. A red-emitting analog of FLASH (named REASH) is also known in the art and may be used for FRET experiment in accordance with this invention by use of, inter alia, YFP; see Galetta, Science 296 (2002), 503-507.

The detection portions/labels present in the recombinant seven-transmembrane protein of the invention facilitate the detection of a conformational change, which, in turn, is indicative for change of the energy emitted by the detection portions/detectable labels.

In one embodiment of the recombinant seven-transmembrane receptor of the invention, these detection labels are portions of a split fluorescent protein. Preferably, this split fluorescent portions is a split green fluorescent protein (split GFP). The term "green fluorescent protein" or "GFP" as used throughout the present application refers to the GFP initially cloned by Prasher (Gene 111 (1992), 229-233) from *Aequorea victoria* and mutants thereof showing GFP activity. The term "GFP activity" refers to the known properties of a GFP, i.e. fluorescence emission upon excitation by a suitable light, the capacity of autocatalytic maturation involving folding into tertiary structure and the formation of the chromophore and the independence of any co-factors or metabolic energy supply for carrying out fluorescence as well as autocatalytic maturation. These properties are well known in the art and for example

reviewed by Tsien (Ann. Rev. Biochem. 67 (1998), 509-544). For the purposes of the present invention, unless otherwise stated, any detectable emission wavelength of a GFP mutant can be useful for applying the recombinant seven-transmembrane protein of the invention. In the prior art, many GFP mutants are described, wherein specific amino acid residues are substituted with the effect of an improved fluorescence efficiency and/or a shifted excitation and/or emission wavelength (see, e.g., Helm, Methods Enzymol. 302 (1999), 408-423; Heikal et al., PNAS 97 (2000), 11996-12001). Particularly, mutating glutamine in position 69 to methionine can reduce the inherent pH and halide sensitivity of eYFP (Griesbeck et al., J. Biol. Chem. (2001) 276, 29188-29194). Thus, if eYFP, or a derivative thereof having substantially the same excitation and emission spectrum, is used as one detection portion of the fusion protein of the invention, it is preferred that the eYFP or derivative thereof shows this mutation. Yet, as shown in the appended examples, YFP is also useful in accordance with this invention. Examples for GFP mutants useful for applying the invention include (enhanced) yellow fluorescent protein ((e)YFP), (enhanced) cyan fluorescent protein ((e)CFP), (enhanced) blue fluorescent protein ((e)BFP), (enhanced) green fluorescent protein ((e)GFP), DsRED, Citrine and Sapphire. Within the scope of the present invention, any GFP mutant or functional analog of GFP may be used as long as it shows fluorescent activity. Preferably, such GFP variants/mutants are encoded by a nucleic acid molecule that hybridizes, preferably under stringent conditions, with the nucleotide sequence encoding the wild-type GFP, or with variants/mutants as the sequence depicted under SEQ ID NOS: 7 to 10. These GFP-mutants/variants shown in SEQ ID NOS: 7 to 10 relate to the most preferred GFP variants to be employed in this invention, namely enhanced cyan fluorescent protein (eCFP) and yellow fluorescent protein (YFP). Suitable preferred hybridization conditions and sequence identity values for preferred hybridizing nucleotide sequences encoding a mutant GFP are mentioned below in connection with functional analogs of the recombinant seven-transmembrane protein of the invention.

The term "split fluorescent protein" refers to a fluorescent protein the amino acid sequence of which is divided into two portions, whereby upon secondary spatial joining of these portions, the split fluorescent protein assumes a three-dimensional



structure which allows it to emit fluorescence when excited by light of a suitable wavelength. It is for example contemplated that the split fluorescent protein is a split GFP, as it has been described by Baird (Proc. Natl. Acad. Sci. USA 96 (1999), 11241-11246). Following the teachings of the prior art, it is possible for a person skilled in the art to divide a GFP into two split GFP portions for fusing them to the 1<sup>st</sup> and 3<sup>rd</sup> loop or a C-terminus of the recombinant seven-transmembrane protein of the invention. It is furthermore conceivable that other fluorescent proteins than GFP, e.g. those mentioned infra, may be split so as to constitute two detection portions in the same manner as split GFP described herein.

In another embodiment of the present invention, the first detection label is an energy-emitting protein portion and the second detection portion is a fluorescent protein label or vice versa. In connection with this embodiment, it is unimportant on which part of the seven-transmembrane protein the first detection portion is located with respect to the other part defined herein, i.e. whether said detection label is located on the first or third intracellular loop or the C-terminus of the inventive recombinant protein. The term "energy-emitting protein portion" refers to proteins capable of radiative energy emission which can (i) take up energy in a suitable form and (ii) transmit at least part of this energy by resonance energy transfer (RET) to the second detection label being a fluorescent protein portion which is thereby elicited to energy emission. The form of energy uptake may be anything that is conceivable to the person skilled in the art and may involve, e.g., a chemical reaction (chemiluminescence or bioluminescence) or absorption of radiation (fluorescence or phosphorescence).

The term "fluorescent protein portion" refers to proteins that are capable of fluorescence, i.e. to absorb energy from radiation of a certain wave length, e.g. ultra-violet or visible light, and to emit this energy or a part thereof by radiation, wherein the emitted radiation has a higher wavelength than the eliciting radiation. There are many examples of fluorescent proteins described in the literature that may be useful in connection with the present invention such as GFPs as mentioned above, fluorescent proteins from non-bioluminescent organisms of the class Anthozoa (WO 00/34318, WO 00/34319, WO 00/34320, WO 00/34321, WO 00/34322, WO 00/34323, WO 00/34324, WO 00/34325, WO 00/34326, WO

00/34526) or the fluorescent protein bmFP from *Photobacterium phosphoreum* (Karatani, Photochem. Photobiol. 71 (2000), 230). Preferred, however, are fluorescent proteins being a YFP and eCFP as employed in the appended examples.

The term "resonance energy transfer" (RET) refers to a non-radiative transfer of excitation energy from a donor (first detection portion) to an acceptor molecule (second detection portion). The conformational change of the recombinant seven-transmembrane receptor results in a detectable change of RET between the detection portions. Such a change can for instance be taken from a comparison of the emission spectra of a recombinant seven-transmembrane receptor in the absence of a suitable binding compound/ligand/agonist or antagonist with the same recombinant seven-transmembrane receptor in the presence of such a compound. If, for example, RET is increased, the emission peak of the acceptor is raised and the emission peak of the donor is diminished. Thus, the ratio of the emission intensity of the acceptor to that of the donor is indicative for the degree of RET between the detection portions. The conformational change of the recombinant seven-transmembrane protein upon binding of a compound, ligand, agonist or antagonist may result either in a decrease or an increase of the distance between the detection portions.

In a most preferred embodiment of the invention, recombinant membrane receptor of the invention, is a G-protein-coupled receptor comprising at least two labels is selected from the group consisting of:

- (a) a polypeptide as shown in SEQ ID NOS: 12, 14 or 16;
- (b) a polypeptide encoded by a nucleic acid sequence as depicted in any one of SEQ ID NOS: 11, 13 or 15;
- (c) a recombinant membrane receptor as defined herein encoded by a nucleotide sequence which hybridizes to a nucleotide sequence as defined (b); and
- (d) a recombinant membrane receptor as defined herein encoded by a nucleic acid sequence being degenerate as a result of the genetic code to a nucleic acid sequence as defined in (b) or (c).

In accordance with this invention, a recombinant membrane receptor is described, wherein the third intracellular loop is or comprises the first detectable label whereby said third intracellular loop is selected from the group consisting of

- (a) a polypeptide depicted in SEQ ID NOS: 18, 22 or 26;
- (b) a polypeptide encoded by a nucleic acid sequence as depicted in SEQ ID NOS: 17, 21 or 25;
- (c) a third intracellular loop encoded by a nucleotide sequence which hybridizes to a nucleotide sequence as defined (b); and
- (d) a third intracellular loop encoded by a nucleic acid sequence being degenerate as a result of the genetic code to a nucleic acid sequence as defined in (b) or (c).

Accordingly, another embodiment of the present invention relates to nucleic acid molecules comprising a nucleotide sequence encoding the recombinant seven-transmembrane protein of the present invention.

The term "nucleic acid molecule" means DNA or RNA or both in combination or any modification thereof that is known in the state of the art (see, e.g., US 5525711, US 4711955, US 5792608 or EP 302175 for examples of modifications). Such nucleic acid molecule(s) are single- or double-stranded, linear or circular and without any size limitation. The nucleic acid molecules of the invention can be obtained for instance from natural sources or may be produced synthetically or by recombinant techniques, such as PCR. In a preferred embodiment, the nucleic acid molecules of the invention are DNA molecules, in particular genomic DNA or cDNA, or RNA molecules. Preferably, the nucleic acid molecule is double-stranded DNA. Particular inventive nucleic acid molecules are depicted in SEQ ID NOS: 11, 13 and 15.

The nucleic acid molecule comprising a nucleotide sequence encoding it is a recombinant nucleic acid molecule, i.e. a nucleic acid molecule that has been produced by a technique useful for artificially combining nucleic acid molecules or parts thereof that were beforehand not connected as in the resulting recombinant nucleic acid molecule. Suitable techniques are for example available from the

prior art, as represented by Sambrook and Russell (2001), *Molecular Cloning: A Laboratory Manual*, CSH Press and Ausubel, *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y. (1989) as well as Vilardaga (1995), *Biotechniques* 18, 605-606. Furthermore, the corresponding techniques are illustrated in the appended examples. Said techniques comprise in particular site-directed mutagenesis.

In this context the term "hybridization" means hybridization under conventional hybridization conditions. They may be low stringent, preferably stringent (i.e. high stringent) hybridization conditions, as for instance described in Sambrook et al., *Molecular Cloning, A Laboratory Manual*, loc. cit. In an especially preferred embodiment the term "hybridization" means that hybridization occurs under the following conditions.

Furthermore, the present invention relates to expression cassettes comprising the above-described nucleic acid molecule of the invention and operably linked thereto control sequences allowing expression in prokaryotic or eukaryotic cells.

Suitable expression control sequences include promoters that are applicable in the target host organism or host cell. Such promoters are well known to the person skilled in the art for diverse hosts from prokaryotic and eukaryotic organisms and are described in the literature. For example, such promoters can be isolated from naturally occurring genes or can be synthetic or chimeric promoters. Likewise, the promoter can already be present in the target genome and will be linked to the nucleic acid molecule by a suitable technique known in the art, such as for example homologous recombination. Specific examples of expression control sequences and sources from where they can be derived are given further below and in the appended examples. All constructs were in pcDNA3.

Expression cassettes according to the invention are particularly meant for an easy to use insertion into target nucleic acid molecules such as vectors or genomic DNA. For this purpose, the expression cassette is preferably provided with nucleotide sequences at its 5'- and 3'-flanks facilitating its removal from and

Insertion into specific sequence positions like, for instance, restriction enzyme recognition sites or target sequences for homologous recombination as, e.g. catalyzed by recombinases.

The present invention also relates to vectors, particularly plasmids, cosmids, viruses and bacteriophages used conventionally in genetic engineering, that comprise a nucleic acid molecule or an expression cassette of the invention.

In a preferred embodiment of the invention, the vectors of the invention are suitable for the transformation of fungal cells, plant cells, cells of microorganisms (i.e. bacteria, protists, yeasts, algae etc.) or animal cells, in particular mammalian cells. Preferably, such vectors are suitable for the transformation of human cells. Methods which are well known to those skilled in the art can be used to construct recombinant vectors; see, for example, the techniques described in Sambrook and Russell (2001), loc. cit.. Alternatively, the vectors may be liposomes into which the nucleic acid molecules or expression cassettes of the invention can be reconstituted for delivery to target cells. Likewise, the term "vector" refers to complexes containing such nucleic acid molecules or expression cassettes which furthermore comprise compounds that are known to facilitate gene transfer into cells such as polycations, cationic peptides and the like.

In addition to the nucleic acid molecule or expression cassette of the invention, the vector may contain further genes such as marker genes which allow for the selection of said vector in a suitable host cell and under suitable conditions. Generally, the vector also contains one or more origins of replication.

Advantageously, the nucleic acid molecules contained in the vectors are operably linked to expression control sequences allowing expression, i.e. ensuring transcription and synthesis of a translatable RNA, in prokaryotic or eukaryotic cells.

In one aspect, the expression of the nucleic acid molecules of the invention in prokaryotic or eukaryotic cells is interesting because it permits a more precise characterization of the function of the recombinant seven-transmembrane protein encoded by these molecules. In addition, it is possible to insert different additional mutations into the nucleic acid molecules by methods usual in molecular biology

(see for instance Sambrook and Russell (2001), loc. cit.), leading to the synthesis of proteins possibly having modified properties, e.g. as concerns binding affinity or energy emission (e.g. RET) efficiency. In this regard, it is possible to mutate the nucleic acid molecules present in the vector by inserting or deleting coding sequences or to introduce amino acid substitutions by replacing the corresponding codon triplets.

For genetic engineering, e.g. in prokaryotic cells, the nucleic acid molecules of the invention or parts of these molecules can be introduced into plasmids which permit mutagenesis or sequence modification by recombination of DNA sequences. Standard methods (see Sambrook and Russell (2001), loc. cit.) allow base exchanges to be performed or natural or synthetic sequences to be added. Similarly, for expression in eukaryotic cells, corresponding expression vectors, like pCDNA3 may be employed. DNA fragments can be connected to each other by applying adapters and linkers to the fragments. Moreover, engineering measures which provide suitable restriction sites or remove surplus DNA or restriction sites can be used. In those cases, in which insertions, deletions or substitutions are possible, in vitro mutagenesis, "primer repair", restriction or ligation can be used. In general, sequence analysis, restriction analysis and other methods of biochemistry and molecular biology are carried out as analysis methods. The expression of the nucleic acid molecule of the present invention is preferably in a stable cell line. Procedure for selection of stably transfected cell lines are known in the art; see, inter alia, Vilardaga (2001), JBC 276, 33435-33443. Preferred host cells are CHO-cells, HEK293 cells, PC12 cells or even primary cells like primary cardiomyocytes or primary brain cells like cultured neurons, cerebral cortex astrocytes, dorsal root ganglia cells and the like.

In a further embodiment, the invention relates to a method for producing cells or hosts capable of expressing the recombinant seven-transmembrane protein/receptor of the invention comprising genetically engineering cells or hosts with an above-described nucleic acid molecule, expression cassette or vector of the invention.

Another embodiment of the invention relates to host cells, in particular prokaryotic or eukaryotic cells, genetically engineered with an above-described nucleic acid molecule, expression cassette or vector of the invention, and to cells descended from such transformed cells and containing a nucleic acid molecule, expression cassette or vector of the invention and to cells obtainable by the above-mentioned method for producing the same. As pointed out below, the invention also relates to non-human transgenic animals comprising nucleic acid sequences encoding the recombinant seven-transmembrane protein/receptor of the invention.

Preferably, host cells are bacterial, fungal, insect, plant or animal host cells. In a preferred embodiment, the host cell is genetically engineered in such a way that it contains the introduced nucleic acid molecule stably integrated into the genome. More preferably the nucleic acid molecule can be expressed so as to lead to the production of the recombinant seven-transmembrane protein of the invention.

A classical overview of different expression systems is for instance contained in *Methods in Enzymology* 153 (1987), 385-516, in Bitter et al. (*Methods in Enzymology* 153 (1987), 516-544) and in Sawers et al. (*Applied Microbiology and Biotechnology* 46 (1996), 1-9), Billman-Jacobe (*Current Opinion in Biotechnology* 7 (1996), 500-4), Hockney (*Trends in Biotechnology* 12 (1994), 456-463), Griffiths et al., (*Methods in Molecular Biology* 75 (1997), 427-440). An overview of yeast expression systems is for instance given by Hensing et al. (*Antoine von Leuwenhoek* 67 (1995), 261-279), Bussineau (*Developments in Biological Standardization* 83 (1994), 13-19), Gellissen et al. (*Antoine van Leuwenhoek* 62 (1992), 79-93), Fleer (*Current Opinion in Biotechnology* 3 (1992), 486-496), Vedvick (*Current Opinion in Biotechnology* 2 (1991), 742-745) and Buckholz (*Bio/Technology* 9 (1991), 1067-1072). Particular preferred expression systems are described in the appended examples and in scientific references cited therein. Expression vectors have been widely described in the literature. As a rule, they contain not only a selection marker gene and a replication origin ensuring replication in the host selected, but also a bacterial or viral promoter and, in most cases, a termination signal for transcription. Between the promoter and the termination signal, there is in general at least one restriction site or a polylinker which enables the insertion of a coding nucleotide sequence. It is possible to use

promoters ensuring constitutive expression of the gene and inducible promoters which permit a deliberate control of the expression of the gene. Bacterial and viral promoter sequences possessing these properties are described in detail in the literature. Regulatory sequences for the expression in microorganisms (for instance *E. coli*, *S. cerevisiae*) are sufficiently described in the literature. Promoters permitting a particularly high expression of a downstream sequence are for instance the T7 promoter (Studier et al., *Methods in Enzymology* 185 (1990), 60-89), lacUV5, trp, trp-lacUV5 (DeBoer et al., in Rodriguez and Chamberlin (Eds), *Promoters, Structure and Function*; Praeger, New York, (1982), 462-481; DeBoer et al., *Proc. Natl. Acad. Sci. USA* (1983), 21-25), lp1, rac (Borbs et al., *Gene* 42 (1986), 97-100). Inducible promoters are preferably used for the synthesis of proteins. These promoters often lead to higher protein yields than do constitutive promoters. In order to obtain an optimum amount of protein, a two-stage process is often used. First, the host cells are cultured under optimum conditions up to a relatively high cell density. In the second step, transcription is induced depending on the type of promoter used. In this regard, a tac promoter is particularly suitable which can be induced by lactose or IPTG (isopropyl- $\beta$ -D-thiogalactopyranoside) (deBoer et al., *Proc. Natl. Acad. Sci. USA* 80 (1983), 21-25). Termination signals for transcription such as the SV40-poly-A site or the tk-poly-A site useful for applications in mammalian cells are also described in the literature. Suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3 (Invitrogen; see also appended examples), pSPORT1 (GIBCO BRL) ) or pCI (Promega).

The transformation of the host cell with a nucleic acid molecule or vector according to the invention can be carried out by standard methods, as for instance described in Sambrook and Russell (2001), loc. cit. The host cell is cultured in nutrient media meeting the requirements of the particular host cell used, in particular in respect of the pH value, temperature, salt concentration, aeration, antibiotics, vitamins, trace elements etc. The recombinant seven-transmembrane protein according to the present invention can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol



precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. In view of the possession of seven-transmembrane regions, the recombinant protein may be purified applying detergents, like TritonX-100 or SDS. Protein refolding steps can be used, as necessary, in completing the configuration of the protein. Such a purified recombinant protein may, inter alia, be reassembled and/or introduced into artificial biological membrane, like liposomes, crude membrane preparations or lipid bilayers.

Accordingly, a further embodiment of the invention relates to a method for producing the recombinant seven-transmembrane protein of the invention comprising culturing the above-described host cells under conditions allowing the expression of said recombinant protein and recovering said recombinant protein from the membranes of the host cell or host organism. Since the recombinant protein is localized in the membranes of the host cells, the protein can be recovered from the cultured cells by detergent-treatment.

Moreover, the invention relates to recombinant seven-transmembrane proteins which are obtainable by a method for their production as described above.

The recombinant seven-transmembrane protein of the present invention may, e.g., be a product of chemical synthetic procedures and is preferably produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect or mammalian cells in culture). Since the inventive protein is a membrane protein, it is preferably produced in a eukaryotic host cell or organism. Depending upon the host employed in a recombinant production procedure, the expressed protein may be modified, e.g. glycosylated or may be non-glycosylated, phosphorylated, palmytolated, ubiquitinated, methylated and the like. The recombinant protein of the invention may also include an initial methionine amino acid residue. The protein according to the invention may be further modified to contain additional chemical moieties not normally part of the protein. Those derivatized moieties may, e.g., improve the stability, solubility, the

biological half life or absorption of the protein. The moieties may also reduce or eliminate any undesirable side effects of the protein and the like. An overview for these moieties can be found, e.g., in Remington's Pharmaceutical Sciences (18<sup>th</sup> edition, Mack Publishing Co., Easton, PA (1990)).

The present invention furthermore relates to non-human transgenic organisms, i.e. multicellular organisms comprising a nucleic acid molecule encoding a fusion protein of the invention or an expression cassette or vector as described above, preferably stably integrated into its genome, at least in a subset of the cells of that organism, or to parts thereof such as tissues or organs. Most preferably, such non-human transgene origin is a mammal like mouse, a rat, a sheep, a goat, a pig, a dog, a rat or a horse.

The transgenic animal expressing the recombinant seven-transmembrane protein of the present invention are particularly useful in pharmacological studies, screening and identification method as provided herein. It is of note that in particular for these studies not only cells but also organs or parts of organs of said non-human transgenic animals are particularly useful. It is envisaged that, for example brains or slice cultures of brain of the herein described non-human transgenic animal are employed in the screening and identification method provided herein. Besides the non-human transgenic animals which are mammals, it is also envisaged that said non-human transgenic organisms may be an amphibian, an insect, a fungi or even a plant. Particular preferred non-human transgenic animals in this context are *Drosophila*, *C. elegans*, *Xenopus* as well as yeasts like *S. pombe* or *S. cerevisiae* or the *Aspergillus* species. Transgenic plants comprise, but are not limited to, wheat, tobacco, parsley or *Arabidopsis*.

As mentioned herein above and as in particular illustrated in the appended examples, the recombinant seven-transmembrane proteins defined herein are in particular useful in screening and identification methods for molecules or compounds which are capable of modifying the biological and/or pharmacological action of seven-transmembrane proteins, in particular of GPCRs and proto-

oncogenes. As the examples show, the present invention is based on the surprising finding that intramolecular RET-analysis can be carried out on rather complex proteins comprising seven-transmembrane regions. It was in particularly surprising that recombinant seven-transmembrane proteins as defined herein provide for a direct measurement system to investigate activation (or deactivation) of GPCR in very fast time frames. The prior art, like Gether (1995), JBC 270, 28268-28275; Jensen (2001), JBC 276, 9279-9290, Ghanouni (2001), JBC 276, 24433-24436 or Ghanouni (2001), PNAS 98, 5997-6002 has provided for screening methods comprising spectroscopic studies where, inter alia, agonists-mediated changes in GPCRs could be measured on a minute time scale. This is much slower than the biological responses to receptor activation which can occur within seconds, even milliseconds. Accordingly, the present invention provides for the first time means and methods whereby activation (as well as deactivation) of seven-transmembrane proteins, in particular GPCRs may be observed with a high resolution and within physiological kinetics. In particular, high resolution assays for conformational changes/switches of receptor activation in living cells are provided.

Accordingly, the present invention provides in one embodiment a method for identifying molecules or compounds which are capable of activating, deactivating or inactivating the (biological/pharmacological) function of (a) seven-transmembrane receptor(s), comprising the steps of

- (a) contacting the recombinant membrane receptor, a host or a host cell as defined herein with (a) molecule(s) or compound(s) to be tested; and
- (b) measuring as a response whether said molecule(s) or compound(s) to be tested lead(s) to a modification of a signal provided by said at least two detectable labels.

Furthermore, a method of screening for molecules or compounds which are activators (agonists) or inhibitors (antagonists) of the (biological/pharmacological) function of (a) seven-transmembrane receptor(s) is provided, said method comprising the steps of

- (a) contacting a recombinant membrane receptor, a host or a host cell as defined herein with the molecule or compound to be tested;
- (b) measuring and/or detecting a response comprising a modification of a signal provided by said at least two detectable labels; and
- (c1) comparing said response to a standard response as measured in the absence of said candidate molecule/compound;
- (c2) comparing said response to the response of a control membrane receptor which comprises at least two detectable labels on the C-terminus; or
- (c3) comparing said response to control seven-transmembrane receptor which comprises only one detectable label.

Similarly, the invention provides for a method for identifying molecules or compounds which are capable of eliciting a (biological/pharmacological) response of (a) seven-transmembrane protein(s), comprising the steps of

- (a) contacting a recombinant membrane protein, a host or host cell of the invention with the molecule or compound to be tested; and
- (b) identifying among these molecules/compounds the molecules/compounds which are capable of eliciting a change in energy emitted by said at least two detectable labels comprised on the recombinant membrane receptor as defined above.

According to the methods provided herein, the invention provides for identifying, characterizing, screening as well as derivatized molecules which are capable of interacting with seven-transmembrane protein in particular with GPCRs whereby said interaction may lead to an activation, a partial activation, an inhibition or a partial inhibition of the biological and/or pharmacological function of said seven-transmembrane protein. Therefore, the present invention provides for distinct screening as well as identification methods for agonists, partial agonists, inverse agonists as well as antagonists of seven transmembrane receptors, in particular of GPCRs. In context of this invention as well as in accordance with the pharmacological sciences, the term "agonist" can be confined as a molecule or a compound that binds to and activates the corresponding receptor. As "partial

agonists" the art defines molecules/compounds that behave like agonists, but that, even at high concentrations, cannot activate the receptors to the same maximal extend as full agonists. The term "inverse agonist" relates to molecules/compounds that bind to and inhibit activity of the corresponding receptor. These inverse agonists are of particular importance and visible, when the receptors exhibit intrinsic agonist-independent activity. The term "antagonist" relates to molecules/compounds that bind to receptors but do not alter the intrinsic activity of said receptor. They may also prevent binding of the corresponding receptor ligand and they may prevent the binding and activation of the receptors by their agonists or partial agonists.

In accordance with the present invention, the term "antagonist" denotes molecules/substances, which are capable of inhibiting and/or reducing an agonistic effect. The term "antagonist" comprises competitive, non-competitive, functional and chemical antagonists as described, inter alia, in Mutschler, "Arzneimittelwirkungen" (1986), Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, Germany. The term "partial antagonist" in accordance with the present invention means a molecule/substance that is capable of incompletely blocking the action of agonists through, inter alia, a non-competitive mechanism. As "agonist", in accordance with this invention, molecules/substances are denoted which have an affinity as well as an intrinsic activity. Mostly, said intrinsic activity ( $\alpha$ ) is defined as being proportional to the quotient of the effect, triggered by said agonist ( $E_A$ ) and the effect which can be maximally obtained in a given biological system ( $E_{max}$ ): therefore, the intrinsic activity can be defined as

$$\alpha \sim \frac{E_A}{E_{max}}$$

The highest relative intrinsic activity results from  $E_A/E_{max}=1$ . Agonists with an intrinsic activity of 1 are full agonists, whereas substances/molecules with an intrinsic activity of  $>0$  and  $<1$  are partial agonists. Partial agonists show a dualistic effect, i.e. they comprise agonistic as well as antagonistic effects.

The person skilled in the art can, therefore, easily employ the compounds and the methods of this invention in order to elucidate the agonistic and/or antagonistic effects and/or characteristics of a compound/molecule/substance to be identified and/or characterized in accordance with any of the above described methods.

The identification and/or characterization of molecules which are capable of interacting with seven-transmembrane receptors, may be, inter alia, achieved by transfecting an appropriate host with a nucleic acid molecule encoding the same and as defined above. Said hosts comprise, but are not limited to, HEK 293 cells, CHO-cells, (primary) cardiomyocytes, (primary) cultured nerve cells, or frog oocytes. After expression of a recombinant seven-transmembrane protein of the invention, membrane currents may be deduced in the absence and/or presence of the molecule to be identified and/or characterized. Methods for the deduction of membrane currents are well known in the art and comprise, e.g., patch clamp methods as described in Hamill, Pflüger's Arch. 391 (1981), 85-100 or two-electrode voltage clamp in oocytes, as described in Methfessel, Pflügers Archive 407 (1986) 577-588.

However, as the appended examples illustrate, the particular preferred measurement methods comprise the FRET- or BRET-measurements as will be detailed below. Yet, also GIRK current measurements are envisaged and shown in the appended examples.

Furthermore, the present invention relates to a method of screening for molecules which are capable of interacting with seven-transmembrane proteins, comprising the steps of (a) contacting a recombinant seven-transmembrane protein of the invention or as encoded by a nucleic acid molecule, a vector or a host of the invention with a candidate molecule; and (b) measuring and/or detecting a response; and (c) comparing said response to a standard response as measured in the absence of said candidate molecule.

Potential candidate molecules or candidate mixtures of molecules may be, inter alia, substances, compounds or compositions which are of chemical or biological origin, which are naturally occurring and/or which are synthetically, recombinantly and/or chemically produced. Thus, candidate molecules may be proteins, protein-fragments, peptides, amino acids and/or derivatives thereof or other compounds, such as ions, which bind to and/or interact with, inter alia, GPCRs.

A person skilled in the art will immediately appreciate that the methods of the invention may present an important contribution to pharmacological research, in particular in the field of drug screening. Thus, corresponding techniques for drug screening described in the literature are incorporated herein by reference. This includes for instance Kyranos (Curr. Opin. Drug. Discov. Devel. 4 (2001), 719-728), Pochapsky (Curr. Top. Med. Chem. 1 (2001), 427-441) and Bohets (Curr. Top. Med. Chem. 1 (2001), 367-383).

According to the present embodiment, in principle any kind of cell, membrane, membrane preparation or liposome may be used for the present method that is amenable to optical detection. The cell to be used can be transformed so as to express a heterologous protein, i.e. the recombinant seven-transmembrane protein of the present invention. Thus, the cells may be single cells such as bacteria, yeasts, protozoa or cultured cells, e.g., of vertebrate, preferably mammalian, more preferably human origin. For certain applications, it may be useful to take pathogenetically affected cells such as tumor cells or cells infected by an infectious agent, e.g. a virus, wherein preferentially measurements are conducted in comparison with corresponding healthy cells. Likewise, the cells may be part of a tissue, organ or organism, in particular of a non-human transgenic animal defined above.

The candidate compounds or test compounds can in principle be taken from any source. They may be naturally occurring substances, modified naturally occurring substance, chemically synthesized substances or substances produced by a

transgenic organism and optionally purified to a certain degree and/or further modified. Practically, the candidate compound may be taken from a compound library as they are routinely applied for screening processes.

The term "contacting" refers to the addition of a candidate compound/test compounds to the analyzed cell in a way that the compound may become effective to the cell at the cell surface or upon cellular uptake. Typically, the candidate compound or a solution containing it may be added to the assay mixture. Step (a) of the methods of the present invention, i.e. the "contacting step" may likewise be accomplished by adding a sample containing said candidate compound or a plurality of candidate compounds to the assay mixture. If such a sample or plurality of compounds is identified by the present method to contain a compound of interest, then it is either possible to isolate the compound from the original sample or to further subdivide the original sample, for example, if it consists of a plurality of different compounds, so as to reduce the number of different substances per sample and repeat the method with the subdivisions of the original sample. Depending on the complexity of the sample, the steps described herein can be performed several times, preferably until the sample identified according to the method of the invention only comprises a limited number of or only one substance(s). Preferably said sample comprises substances of similar chemical and/or physical properties, and most preferably said substances are identical.

Step. (b), i.e. the "measuring or identification step" may be carried out in accordance with the explanations regarding measuring a change in energy emission of the fusion proteins of the invention as given hereinabove. Particularly preferred are optical measurement techniques that allow a resolution of fluorescence on the level of single cells, preferably at the subcellular level. Suitable imaging techniques are described in the literature such as in Periasamy A., *Methods in Cellular Imaging*, 2001, Oxford University Press or in *Fluorescence Imaging Spectroscopy and Microscopy*, 1996, edited by: X.F. Wang; Brian Herman. John Wiley and Sons. They may involve fluorescence, preferably confocal, microscopy, digital image recording, e.g. by way of a CCD camera, and suitable picture analysis software. The appended



examples also provide for useful settings for measuring candidate compounds. Preferentially, step (b) is carried out by running parallel control experiments. For instance, a corresponding cell expressing the same recombinant seven-transmembrane protein may be observed under corresponding conditions as in steps (a) and (b), however, without contacting a candidate compound.

Accordingly, potential candidate molecules may be contacted with a cell, such as an oocyte, a HEK 293 cell, a CHO cell, a PC12 cell, or an (primary) cardiomyocytes which express a recombinant seven-transmembrane protein of the invention or with a membrane patch, a membrane preparation, comprising a recombinant seven-transmembrane protein of the invention and measuring a corresponding response (inter alia, a dose-response, a current-response, or a concentration response) in order to elucidate any effect said candidate molecule causes. Said response is most preferably measured by methods provided herein and in particular by FRET or BRET technology.

Within the scope of the present invention are also methods for identifying, characterizing and for screening of molecules which are capable of interacting with seven-transmembrane receptors which comprise so-called high-throughput screening methods and similar approaches which are known in the art (Spencer, *Biotechnol. Bioeng.* 61 (1998), 61-67; Oldenburg, *Annu. Rep. Med. Chem.* 33 (1998), 301-311; Milligan, *Trends Pharmacol. Sci.* 20 (1999), 118-124) carried out using 96-well, 384-well, 1536-well (and other) commercially available plates. Further methods to be employed in accordance with the present invention comprise, but are not limited to, homogenous fluorescence readouts in high-throughput screenings (as described, inter alia, in Pope, *Drug Discovery Today* 4 (1999), 350-362). The method of the present invention for identification, characterization and/or screening of molecules capable of interacting with seven transmembrane-proteins, in particular GPCRs, can, inter alia, employ hosts as defined herein which express the recombinant protein of the present invention. Cell-based assays, instrumentation for said assays and/or measurements are well-known in the art and described, inter alia, in Gonzalez,

Drug Discovery Today 4 (1999), 431-439 or Ramm, Drug Discovery Today 4 (1999), 401-410.

In this context, it is envisaged that agonists as well as antagonists as defined herein may be verified, measured and/or deduced by the methods provided herein. It is also envisaged that modifications, derivatives and the like of known agonists and/or antagonists are measured and screened by the methods of the present invention. For example, the activity of a derivate/derivative of a known agonist or antagonist of a GPCR may be compared to the activity of the corresponding, non-modified (not-derivatized) agonist/antagonist. As non-limiting examples the following list provides for antagonists/agonists of the exemplified GPCRs  $\alpha_2A$ -adrenergic receptor, PTH-receptor and A2A adenosine receptor. Also provided are pharmaceutical/medical uses of the corresponding agonists/antagonists.

1.  $\alpha_2A$ -adrenergic receptor:

Compounds:

Agonists: clonidine and similar drugs (centrally acting antihypertensive; treatment of withdrawal syndrome, open angle glaucoma, diarrhea in patients with neuropathies); oxymetazoline and many similar drugs (direct vasoconstrictor: decongestion in rhinitis, conjunctivitis etc); apraclonidine, brimonidine (open angle glaucoma); dexmedetomidine (anaesthetic, sedative).

Possible further medical use: psychiatric and neurologic (presynaptic inhibition of neurotransmitter release)

Antagonists: Phentolamine (peripherally acting antihypertensive, particularly in pheochromocytoma)

Possible further medical use: psychiatric (presynaptic desinhibition), functional treatment of degenerative CNS diseases

Diseases (in addition to those mentioned above) to be treated by agonists comprise

Risk factor in cardiovascular diseases (increased release of neurotransmitter if receptor dysfunctional: hypertension, heart failure; in CNS-dysfunctions, agonists and antagonists are envisaged)

## 2. PTH-receptor:

Agonists: parathyroid hormone (PTH) itself and smaller fragments thereof, e.g. PTH(1-34) (hypoparathyroidism, osteoporosis, diagnostic use in different forms of hypoparathyroidism).

Antagonists: Small fragments of PTH, e.g. PTH(7-34)

possible medical use: hyperparathyroidism

Diseases (in addition to those mentioned above) to be treated by agonists comprise diseases of vitamin-D-metabolism, Ca-homeostasis; diseases of defective bone formation or metabolism (like osteoporosis)

## 3. A<sub>2A</sub> adenosine receptor:

Agonists: Adenosine and several of its purine modified derivatives, e.g. N-ethylcarboxamidoadenosine (NECA);

Effects: Vasodilation, inhibition of platelet aggregation, various CNS-effects.

Possible medical use: disorders of platelet aggregation and consequences of such disorders (atherosclerosis, stroke, myocardial infarction, heart failure), vasodilation for hypertension, vasospasm, various psychiatric/neurological illnesses have been postulated as potential indications.

Antagonists: theophylline, caffeine and other methylxanthines

Bronchial asthma - bronchodilation, central stimulation, diuresis, apnea – particularly in infants (not actually proven that this receptor subtype is the target in these therapeutic effects – other homologous adenosine receptors (A<sub>1</sub>, A<sub>2B</sub>, A<sub>3</sub>) might also be involved)

possible medical use: disorders of attention such as e.g. ADHS

Further GPCRs to be modified in form of recombinant seven-transmembrane receptors of the present invention comprise, but are not limited to:  $\beta_1$ -,  $\beta_2$ -,  $\beta_3$ -,

$\alpha_{1A, B, \Delta}$ ,  $\alpha_{2A, B, X}$ -adrenergic receptor;  $M_{1-5}$  muscarinic receptor;  $H_1$ -,  $H_2$ -histamine receptor,  $AT_1$ -angiotensin receptor; serotonin-receptor(s) as well as opiate receptors. Corresponding agonists/antagonists and medical indications are given in the following table. Again it is envisaged that derivatives and/or modifications of the given agonists/antagonists be tested by the methods provided herein.

Receptor	Agonists (Indication)	Antagonists (Indication)
<u>Adrenergic</u> $\beta_1$	Noradrenaline, adrenaline, dobutamine (cardiac shock, acute heart failure)	metoprolol, bisoprolol and many others (hypertension, arrhythmias, heart failure, myocardial infarction)
$\beta_2$	Isoprenaline, terbutaline, salbutamol and many others (bronchial asthma – bronchodilation, premature labour)	propranolol and many others (as for $\beta_1$ ; also: open angle glaucoma, hyperthyroidism, tremor, some forms of anxiety)
$\beta_3$	Isoprenaline, several experimental compounds, e.g. BRL37344 (lipolysis)	several experimental compounds, e.g. ICI118551
$\alpha_{1A, B, \Delta}$	Noradrenaline, adrenaline, phenylephrine (Constriction of smooth muscle, eg. vasoconstriction, activation of gastrointestinal smooth muscles)	prazosin, terazosin, tamsulosin and others (hypertension, prostatism)
$\alpha_{2A, B, X}$	Noradrenaline, adrenaline, clonidine, oxymetazoline	yohimbin (use as aphrodisiac)

	(see above)	
<u>Muscarinic</u> M <sub>1-5</sub>	Acetylcholine, carbachol, pilocarpine (glaucoma, activation of gastrointestinal motility, tacharrhythmias)	Atropine, scopolamine (bradycardia, (pre-) anesthesia, bronchial asthma, vagal reflexes, poisoning with E605 and similar compounds including nerve gas)
<u>Histamine</u> H <sub>1</sub>	histamine	loratadine, clemastine (antiallergic, sedative-hypnotic)
H <sub>2</sub>	histamine	ranitidine and many others (gastric acid secretion: gastric and duodenal ulcers)
<u>Angiotensin</u> AT <sub>1</sub>	angiotensin	losartan (antihypertensive)
<u>Serotonin (5-HT)</u> 14 receptor subtypes, e.g.: 5-HT <sub>1B/D</sub>  5-HT <sub>2C</sub>  5-HT <sub>4</sub>	sumatriptan and others (migraine)   partial agonists: metoclopramide (gastrointestinal motility disorders)	   methysergide, pizotifen (migraine, carcinoid syndrome)
<u>Opiate</u> $\mu$	morphine and many others (analgesia, CNS-stimulation (euphoria) and depression, alterations in smooth muscle contraction: increase-spasm/decrease)	naloxone, naltrexone Treatment of opiate overdose, termination of opiate effects
$\kappa$	ethylketocyclazozine and many others (analgesia, diuresis, central nervous system effects)	naloxone, naltrexone as above
$\delta$	etorphine, deltorphin and others (analgesia, alterations in smooth muscle contraction: decrease)	naloxone, naltrexone as above

Additionally, the present invention relates to a method for the production of a pharmaceutical composition comprising the steps of the method of the invention for identifying, characterizing and/or screening of molecules which are capable of interacting with seven-transmembrane receptors, like GPCRs, and further comprising a step, wherein a derivative of said identified, characterized and/or screened molecule is generated. Such a derivative may be generated by, inter alia, peptidomimetics.

The invention furthermore relates to a method for the production of a pharmaceutical composition comprising the steps of the method of the invention for identifying, characterizing, screening and/or derivatizing of molecules which are capable of interacting with seven-transmembrane receptors, like GPCRs and formulating the molecules identified, characterized, screened and/or derivatized in pharmaceutically acceptable form.

In a preferred embodiment, the response or energy changes to be measured in the methods provided herein correspond to an increase or a decrease of fluorescence resonance energy transfer (FRET).

In FRET, both donor and acceptor, i.e. both detection portions, are fluorescent protein portions and, for measuring FRET, the fusion protein is supplied with energy, i.e. radiation, appropriate for exciting energy emission by the first detection portion.

Accordingly, it is a preferred embodiment of the recombinant seven-transmembrane protein of the present invention, that the first detection label is a fluorescent protein portion.

The efficiency of FRET is dependent on the distance between the two fluorescent partners. The mathematical formula describing FRET is the following:  $E = R_0^6 / (R_0^6 + r^6)$ , where E is the efficiency of FRET, r is the actual distance between the fluorescent partners, and  $R_0$  is the Förster distance at which FRET is 50% of the maximal FRET value which is possible for a given pair of FRET partners.  $R_0$ , which can be determined experimentally, is dependent on the relative orientation

between the fluorescent partners ( $\kappa$ ), refractive index of the media ( $n$ ), integral overlap of the emission of the donor with the excitation of the acceptor partner ( $J(\lambda)$ ), and the quantum yield of the fluorescent donor partner ( $Q_D$ ) ( $R_0^6 = 8.79 \times 10^{-25} [\kappa^2 n^4 Q_D J(\lambda)]$  (in  $\text{cm}^6$ )). In classical FRET based applications the orientation factor  $\kappa^2$  is assumed to equal  $2/3$ , which is the value for donors and acceptors that randomize by rotational diffusion prior to energy transfer (Lakovicz, Principles of Fluorescence spectroscopy, second edition, page 370). Thus, at randomized rotational diffusion, the change in ratio is assumed to be only due to a change in distance between the chromophores. For perpendicular dipoles  $\kappa^2$  is 0.

In accordance with the appended examples, a decrease in FRET-signal can be determined by the following equation:  $r(t) = A \times (1 - e^{-t/\tau})$ , where  $\tau$  is the time constant (s) and  $A$  is the magnitude of the signal. When necessary for calculating  $\tau$ , agonist-independent changes in FRET due to photobleaching were subtracted. In order to apply FRET for detection of agonists, antagonists, partial agonists and partial antagonists as well as inverse agonists, the person skilled in the art is capable of selecting suitable detection labels (defined above) for the seven-transmembrane protein of the invention that show a detectable FRET and a detectable change of FRET upon a conformational change in its structure. Preferably, maximum FRET efficiency is at least 5%, more preferably at least 50% and most preferably 80% of the energy released by the first detection label upon excitation. Additionally, the two detection labels need to have a spectral overlap. The greater the overlap of the emission spectrum of the donor with the absorption spectrum of the acceptor, the higher is the value of  $R_0$ . Acceptors with larger extinction coefficients lead to higher  $R_0$  values. In contrast, the overlap in excitation spectra of both detection portions should be small enough to prevent coexcitation of the acceptor chromophore. Likewise, the spectra of both detection portions should only overlap to an extent that discrimination between the two emission signals is still possible.

As detailed in the appended examples, in a particularly preferred embodiment, the first detection portion is cyan fluorescent protein (CFP) and the second detection portion is enhanced yellow fluorescent protein (eYFP).

It has been shown that CFP and YFP are particularly well suited for the recombinant membrane protein of the present invention since they show an efficient change in FRET. CFP and eYFP are well known in the art and nucleic acid molecules containing corresponding coding sequences are commercially available e.g. from Clontech. Said nucleic acid sequences are also shown in appended SEQ ID NOS: 7 to 10.

In a further preferred embodiment of the present invention the methods provided herein are based on the detection of responses or energy changes which comprise an increase or a decrease of bioluminescent resonance energy transfer (BRET). BRET-technology is very well known in the art and, inter alia, described in Angars, (2000) PNAS 97, 3684-3689; in Mercier (2002), JBC 277, 44925-44931; in Barcock, (2003), JBC 278, 3378-3385 or in WO 99/66324. As pointed out herein above a preferred bioluminescent protein is renilla luciferase but also firefly luciferase may be employed. As a preferred fluorescent protein portion in the recombinant seven-transmembrane receptor of the present invention comprising renilla luciferase as a first detection system, enhanced yellow fluorescent protein or yellow fluorescent protein may be employed.

In accordance with the methods provided herein in a most preferred embodiment, the recombinant seven-transmembrane protein of the present invention is located, respectively inserted, into a biological membrane. Most preferably, said biological membrane is a plasma membrane of a cultured cell or is a membrane in (a) cell(s) of an organ or tissue of a non-human transgenic animal expressing the recombinant seven-transmembrane protein of the present invention. Further embodiments of the remaining and/or identification methods of the present invention are given and illustrated in the appended examples. It is of note that in context of the present invention several control as already briefly discussed herein above may be employed. For example, recombinant seven-transmembrane proteins comprising only one detectable label may be used as controls. Such recombinant protein will not provide for any change in energy emitted or to a detectable response which may be measured. Accordingly, the test molecules or test compounds or samples comprised either alone or in combination such



molecules or compounds may be tested in parallel experiment on recombinant seven-transmembrane proteins of the present invention, capable of eliciting a distinct response upon conformational change and recombinant seven-transmembrane receptors which only comprise one of the above-identified detectable labels and are, accordingly, not capable of eliciting a corresponding signal, in particular of eliciting an resonance energy transfer. Another control protein to be employed in accordance with the method of the present invention is the recombinant seven-transmembrane receptor protein which comprises both detectable labels on the C-terminus. Such a control protein is, inter alia, illustrated in appended SEQ ID NO: 38 and was employed in accordance with this invention in the appended examples. This control molecule leads to strong resonance energy transfer. However, upon testing with corresponding ligands, specific for the seven-transmembrane receptor to be tested no further chain of the RET is expected. For example, the here defined control for a PTH receptor comprising both detectable labels in the carboxy terminal provides for a strong FRET signal which does not change when the corresponding ligand, namely parathyroid hormone is applied.

The invention also provides for a diagnostic composition comprising the recombinant membrane protein, the nucleic acid molecule, the vector, the host cell or the organs or cells of the non-human transgenic animal of the invention. Such a diagnostic composition is particularly useful in the methods of the present invention. Similarly, kits are provided which comprise the compounds of the invention, in particular, the recombinant membrane protein, the nucleic acid molecule, the vector, the host cell or the organs or cells of the non-human transgenic animal of the invention.

Finally, the invention relates to the use of the recombinant membrane protein or the nucleic acid molecule, the vector, the host cell or the organs or cells of the non-human transgenic animal of the invention for the detection of (a) modifier(s) of the biological activity of seven-transmembrane receptors in vivo or in vitro.

The embodiments of the method(s) of the invention apply here *mutatis mutandis*.

The Figures show:

**Figure 1 Part 1:** FRET efficiency and time-resolved changes in the FRET signal of PTHR-cam. (A) Overall transmembrane topology of the GPCR-cam constructs. (B) Fluorescence emission spectra of selected PTHR constructs. Shown are the emission spectra of PTHR-CFP<sub>3-loop</sub> (blue), PTHR-YFP<sub>C-term</sub> (yellow) and PTHR-cam (red) upon excitation at 433 nm. (C) Effects of photobleaching. Emission intensities of YFP (535 nm, yellow) and CFP (480 nm, blue) were recorded simultaneously from single cells expressing PTHR-cam using fluorescence microscopy. Emission intensities were recorded before and after the acceptor fluorophore was photobleached by 5 min exposure to light at 480 nm. (D) Time-resolved changes in the ratio  $F_{535}^*/F_{480}^*$  in single HEK293 cells stably expressing PTHR-cam. Emission intensities of YFP (535 nm, yellow), CFP (480 nm, blue) and the ratio  $F_{535}^*/F_{480}^*$  (red) were recorded simultaneously from single cells. Shown are the changes induced by rapid superfusion with 1  $\mu$ M PTH (arrow). The decrease of the ratio  $F_{535}^*/F_{480}^*$  was fitted by a simple mono-exponential curve giving a time-constant in this experiment of 3.5 s. Changes in the ratio are expressed as % decrease from the initial value at  $t = 0$ s.

**Figure 1, Part 2:** Guanine nucleotide sensitivity of agonist binding at the  $\alpha_{2A}$ AR-cam stably expressed in HEK-293 cells. In the presence of exogenous  $G_o$  protein (at a molar ratio 1:100, receptor: $G_o$ ) the binding affinity for the  $\alpha_{2A}$ AR agonist UK14,304 for membranes containing  $\alpha_{2A}$ AR-cam decreases  $\approx 3$ -fold in presence of 10  $\mu$ M GTP $\gamma$ S ( $K_d = 3.4 \pm 0.8$  nM vs  $K_d = 9.6 \pm 1.1$  nM,  $n=3$ ) reflecting the shift of the receptor population to a lower affinity state.

**Figure 2:** Pharmacological properties of the GPCR-cam constructs.

(A-B) Comparison between the binding and signalling properties of PTHR-cam (A),  $\alpha_{2A}$ -AR-cam (B) and their respective wild type receptor stably expressed in HEK293 cells. The expression level of the receptors were  $1.01 \times 10^6$  and  $0.34 \times 10^6$  receptors/cells for PTHR and PTHR-cam, respectively; 24 and 6 pmol/mg for

$\alpha_{2A}$ AR and  $\alpha_{2A}$ AR-cam, respectively. The data are the means $\pm$ S.E. of at least 4 separate experiments carried out in duplicate. (C) Visualization PTHR-cam stably expressed in HEK293 cells by confocal microscopy.

**Figure 3:** Agonist-induced decrease in FRET signal corresponds to receptor activation. (A) *Intact cell* panel shows the effects of the agonist PTH (1  $\mu$ M) and the antagonist PTH(7-34) (3 $\mu$ M) on the ratio  $F_{535}^*/F_{480}^*$  of PTHR-cam in intact HEK293 cells. *Membranes* panel shows the effects of PTH (1  $\mu$ M) in cell membranes prepared from HEK293 cells stably expressing PTHR-cam. The membranes were measured either without further treatment (left), or after treatment with 6 M urea (right). Bars represent the % decrease in the ratio  $F_{535}^*/F_{480}^*$  upon PTH exposure. (B) Effect of the antagonist phentolamine (10  $\mu$ M) on the FRET signal caused by 10  $\mu$ M noradrenaline (NA) in HEK293 cells stably expressing  $\alpha_{2A}$ AR-cam (n=4). (C) Comparison of the guanine nucleotide sensitivity of the FRET signal (*left* panel) and agonist binding (*right* panel) evoked by sub-saturating concentration of UK14304 in membranes containing  $\alpha_{2A}$ AR-cam in the presence of  $G_o$  proteins (ratio receptor: $G_o$  1:100) with or without GTP $\gamma$ S (10  $\mu$ M). Data are the means $\pm$ S.E. of at least 4 separate experiments.

**Figure 4:** Action of the partial agonist clonidine on  $\alpha_{2A}$ AR-cam. Changes in FRET in response to 10  $\mu$ M noradrenaline (NA) or 10  $\mu$ M clonidine added alone or together were recorded in a single HEK293 cell expressing  $\alpha_{2A}$ AR-cam. The recording is representative of 4 independent experiments.

**Figure 5:** Comparison between the dynamics of receptor activation and desensitization of PTHR-cam. The kinetics of activation of PTHR-cam and of  $\beta$ -arrestin2-YFP binding to the PTHR-CFP<sub>C-term</sub> were measured as changes in the ratio  $F_{535}^*/F_{480}^*$  in single cells expressing PTHR-cam or co-expressing PTHR-CFP<sub>C-term</sub> and  $\beta$ -arrestin2-YFP in response to 100 nM PTH. The recordings are expressed as % of the respective maximal response and are representative of at least 3 independent experiments. Note that in the case of PTHR-CFP<sub>C-term</sub>/ $\beta$ -arrestin2-YFP the ratio  $F_{535}^*/F_{480}^*$  does indeed increase, while in the PTHR-cam

the ratio decreases and is depicted as a positive signal just to facilitate the comparison of the kinetics. The *inset* represents a time scale expansion to illustrate the differences in response delays.

**Figure 6:** Dynamics of agonist-mediated receptor conformational change. **(A)** Time-resolved changes in the ratio  $F_{535}^*/F_{480}^*$  of the PTHR-cam (*left* panel) and  $\alpha_{2A}$ AR-cam (*right* panel) expressed in HEK293 cells at various concentrations of PTH and norepinephrine, respectively. **(B)** Relationship between the apparent rate constant,  $k_{obs}$  and agonist concentration.  $k_{obs}$ -values were also obtained from fitting the kinetic data of Figure 6A with a mono-exponential equation. At low concentrations of agonist,  $k_{obs}$ -values were directly proportional to the agonist-concentration, whereas at higher concentrations of agonist the values approached a maximal value of about  $1\text{ s}^{-1}$  and  $26\text{ s}^{-1}$  for PTHR-cam and  $\alpha_{2A}$ AR-cam, respectively. Note that the  $k_{obs}$  values depicted in Fig. 6B saturated at much higher ligand concentrations than the binding data shown in Fig. 2. This is due to the fact that the  $k_{obs}$  values are not measured at equilibrium. Data indicate the mean $\pm$ SEM of at least 7 separate experiments.

The examples illustrate the invention.

#### **Example I: Experimental protocol and sequences encoding membrane receptors of the invention**

**Molecular biology and cell culture.** Site-directed mutagenesis was performed on the human PTH/PTHrP receptor (PTHR) and the mouse  $\alpha_{2A}$ -adrenergic receptor cDNAs. The cDNAs encoding the enhanced yellow- and cyan fluorescent protein (YFP and CFP) were fused to position G418 of the COOH-terminus of the PTHR and inserted between Gly395 and Arg396 into the third intracellular loop of the PTHR, respectively. For the  $\alpha_{2A}$ -adrenergic receptor, YFP was inserted in the third intracellular loop between Ala250 and Ser371, and CFP was fused to Val461 in the C-terminus. Constructions were performed by polymerase chain reactions as described (Villardaga, Biotechniques 18 (1995), 605-606).  $\beta$ -Arrestin2-YFP was

constructed following the procedure described for the construction of arrestin3-GFP (Groarke, J. Biol. Chem. 274 (1999), 23263-23269). Constructions were verified by sequencing. Receptor cDNAs were cloned into pCDNA3 (Invitrogen) for transient and stable expression in mammalian cells. HEK293, CHO and PC12 cell lines served as the expression systems for the wild-type and chimeric receptors. The procedure for the selection of stable cell line has been previously described (Vilardaga, J. Biol. Chem. 276 (2001), 33435-33443).

**Pharmacology.** Ligand binding, receptor number determination, cAMP assays, measurement of  $\alpha_{2A}$ -adrenergic receptor-activated GIRK currents and reconstitution of receptor- $G_o$  coupling were measured as previously described (Vilardaga, (2001) loc. cit.; Bünemann, J. Biol. Chem. 276 (2001), 47512-47517; Vilardaga, J. Biol. Chem. 277 (2002), 8121-8129; Richardson, J. Biol. Chem. 274 (1999), 13525-13533). Saturation and competition binding studies were analysed with the program Prism to calculate  $K_D$ - and  $K_i$ -values.

**Electrophysiology.** Whole cell GIRK currents were measured in HEK293 cells stably expressing 6 pmol/mg membrane protein  $\alpha_{2A}$ -AR-cam 20-28h after transient transfection with GIRK1 and GIRK4 as described previously (Bünemann, (2001), loc. cit.). Membrane currents were recorded using an EPC 9 amplifier and Pulse software (HEKA Instruments) for voltage control, data acquisition and data evaluation. Experimental conditions such as patch pipettes, internal and external solutions, voltage-clamp protocol as well as the superfusion system were the same as described previously (Bünemann, (2001), loc. cit.).

**Fluorescence measurements.** Cells were washed with PBS, scraped from the plate and resuspended in buffer A (137 mM NaCl, 5 mM KCl, 1 mM  $CaCl_2$ , 1 mM  $MgCl_2$ , 20 mM Hepes, 0.1% BSA, pH 7.4) at a density of  $\approx 10^7$  cells/ml. Steady-state fluorescence emission spectra of the cells suspension were measured with a spectrofluorimeter (Perkin-Elmer) in cuvettes containing HEK293 cells expressing the indicated receptors, and were normalized to the respective maxima for PTHR-

CFP<sub>3-loop</sub> and PTHR-cam. PTHR-YFP<sub>C-term</sub> was normalized relative to the maximal response upon exposure to 480 nm light.

**FRET measurements.** Cells grown on coverslips were maintained in buffer A at room temperature<sup>7</sup> and placed on a Zeiss inverted microscope (Axiovert135) equipped with an oil immersion 63x objective and a dual emission photometric system (Till Photonics). Samples were excited with light from a polychrome IV (Till Photonics). In order to minimize photobleaching, the illumination time was set to 5 ms applied with a frequency between 1 and 75 Hz dependent on agonist concentration. FRET was monitored as the emission ratio of YFP to CFP,  $F_{535}/F_{480}$ , where  $F_{535}$  and  $F_{480}$  are the emission intensities at  $535 \pm 15$  nm and  $480 \pm 20$  nm (beam splitter DCLP 505nm) upon excitation at  $436 \pm 10$  nm (beam splitter DCLP 460 nm). The emission ratio was corrected by the respective spill-over of CFP into the 535 nm channel (spill-over of YFP into the 480 nm channel was negligible) to give a corrected ratio  $F_{535}^*/F_{480}^*$ . FRET between CFP and YFP in cells stably expressing the receptor constructs was also determined by donor recovery after acceptor bleaching. The increase in emission at 480nm was  $50 \pm 2\%$  after  $>80\%$  bleaching of YFP (induced by 3-5 min continuous illumination with  $480 \pm 15$ nm). To determine agonist-induced changes in FRET, cells were continuously superfused with buffer A and agonist was applied using a computer assisted solenoid valve controlled rapid superfusion device ALA-VM8, ALA Scientific Instruments (solution exchange 5 to 10 ms). Signals detected by avalanche photodiodes were digitalized using a AD converter (Digidata1322A, Axon Instruments) and stored on PC using Clampex 8.1 software (Axon Instruments). The decrease in FRET ratio was fitted to the equation:  $r(t) = A \times (1 - e^{-t/\tau})$ , where  $\tau$  is the time constant (s) and A is the magnitude of the signal. When necessary for calculating  $\tau$ , agonist-independent changes in FRET due to photobleaching were subtracted.

**Membrane preparation.** Membrane fractions were obtained after two centrifugations at 4°C, first at 800xg for 10 min and the second at 100,000xg for 30 min. Membranes were treated with 6 M urea in 20 mM Hepes, pH 7.4, for 30

min on ice followed by centrifugation at 100,000xg for 15 min. After two washing steps at 4°C in buffer B the membranes were resuspended in a buffer A (without BSA) and immediately used in the experiments. 20 µl aliquots of membranes were subjected to fluorescence microscopy similar to intact cells using a 20x objective.

**The following sequences and constructs have been employed in course of the following experiments:**

**SEQ ID NO: 1: wildtype Mouse alpha2a adenergetic receptor cDNA sequence:**  
*Genbank M99377 (corresponding to human alpha 2A:Genbank: NM\_000681)*

ATGGGCTACCCATACGACGTCCCAGACTACGCCAGCATGGGCTCACTGCAG  
 CCGGATGCCGGCAACAGCAGCTGGAACGGGACCGAAGCGCCCGGAGGCG  
 GCACCCGAGCCACCCCTTACTCCCTGCAGGTGACACTGACGCTGGTTTGCC  
 TGGCTGGCCTGCTCATGCTGTTACAGTATTTGGCAACGTGCTGGTTATTAT  
 CGCGGTGTTACACAGTCGCGCGCTCAAAGCTCCCCAAAACCTCTTCCTGGT  
 GTCCCTGGCCTCAGCGGACATCCTGGTGGCCACGCTGGTCATTCCCTTTTCT  
 TTGGCCAACGAGGTTATGGGTTACTGGTACTTTGGTAAGGTGTGGTGTGAGA  
 TCTATTTGGCTCTCGACGTGCTCTTTTGCACGTCGTCCATAGTGCACCTGTG  
 CGCCATCAGCCTTGACCGCTACTGGTCCATCACGCAGGCCATCGAGTACAA  
 CCTGAAGCGCACGCCGCGTCGCATCAAGGCCATCATTGTCACCGTGTGGGT  
 CATCTCGGCTGTCTCTCCTTCCCGCCACTCATCTCCATAGAGAAGAAGACC  
 AGAAGTGGTATGTCATCTCCTCGTCCATCGGTTCTTCTTCGCGCCTTGCCT  
 CATCATGATCCTGGTCTACGTGCGTATTTACCAGATCGCCAAGCGTCGCACC  
 CGCGTGCCCTCCAGCCGCCGGGGTCCGGACGCCTGTTCCGCGCCGCGCGGG  
 GGGCGCCGATCGCAGGCCCAACGGGCTGGGCCCCGAGCGCGGCGCGGGT  
 CCCACGGGCGCTGAGGCGGAGCCGCTGCCCACCCAGCTTAACGGTGCCCC  
 GGGGGAGCCCCGCGCCCGCCGGGGCCCCGCGATGGGGATGCGCTGGACCTA  
 GAGGAGAGTTTCGTGTCGTCGAGCACGCCGAGCGGCCCCCGGGGCCCCGCGAG  
 ACCCGACCGCGGCCCCCCGAGCCAAGGGCAAGACCCGGGCGAGTCAGGTGA  
 AGCCGGGGGACAGTCTGCCGCGGCGCGGGCCCCGGGGCCGCGGGGGCCGG  
 GGGCTTCGGGGTCCGGGACGAGAGGAGCGCGGCGGGGGGCGCCAAAGC

GTCGCGCTGGCGCGGGAGGCAAAACCGGGAGAAACGCTTCACGTTCTGTGC  
 TGGCGGTGGTGATCGGCGTGTTCTGTGGTGTGTTGGTTTCCGTTCTTTTTCAC  
 CTACACGCTCATAGCGGTGCGGCTGCCCCGGTGCCCAGCCAGCTCTTCAACTT  
 CTTCTTCTGGTTCGGCTACTGCAACAGCTCGCTGAACCCTGTTATCTACACC  
 ATCTTCAACCACGACTTCCGACGCGCCTTCAAGAAGATCCTCTGCCGTGGGG  
 ACAGAAAACGCATCGTGTGA TTC AAC CAC GAC TTC CGA CGC GCC TTC  
 AAG AAG ATC CTC TGC CGT GGG GAC AGA AAA CGC ATC GTG TGA

**SEQ ID NO: 2: wildtype Mouse alpha2a adenergic receptor amino acid sequence:**

MGYPYDVPDYASMGSLQPDAGNSSWNGTEAPGGGTRATPYSLQVTLTLVCLA  
 GLLMLFTVFGNVLVIIAVFTSRALKAPQNLFLVSLASADILVATLVIPFSLANEVMG  
 YWYFGK VWCEIYLALDVL FCTSSIVHLCAISLD RYWSITQAIEYNLKRTPRRIKAIIV  
 TVWVISAVISFPPLISIEKKGAGGGQQPAEPSCKINDQKWYVISSSIGSFFAPCLIM  
 ILVYVRIYQIAKRRTVPSPRRGPDACSAPPGGADRRPNGLGPERGAGPTGAEA  
 EPLPTQLNGAPGEPAPAGPRDGDALDLEESSSEHAERPPGPRRPDRGPRAKG  
 KTRASQVKPGDSLPRRGPGAAGPGASGSGHGEERGGAASRWRGRQNREK  
 RFTFVLAVVIGVFVVCWFPPFFFTYTLIAVGCPVPSQLFNFFFWFGYCNSSLNPVIY  
 TIFNHDFRRFAFKKILCRGDRKRIV

**SEQ ID NO: 3: wildtype Human PTH/PTHrP receptor cDNA sequence:**  
*Genbank: U22401 (corresponding to mouse PTH/PTHrP receptor: NM\_011199)*

ATGGGGACCGCCCGGATCGCACCCGGCCTGGCGCTCCTGCTCTGCTGCCC  
 CGTGCTCAGCTCCGCGTACGCGCTGGTGGA  
 TGCAGATGACGTCATGACTAAAGAGGAACAGATCTTCCTGCTGCACCGTGCT  
 CAGGCCCAGTGCGAAAAACGGCTCAAGG  
 AGGTCCTGCAGAGGCCAGCCAGCATAATGGAATCAGACAAGGGATGGACAT  
 CTGCGTCCACATCAGGGAAGCCCAGGAAA  
 GATAAGGCATCTGGGAAGCTCTACCCTGAGTCTGAGGAGGACAAGGAGGCA  
 CCCACTGGCAGCAGGTACCGAGGGGCGCCC



CTGTCTGCCGGAATGGGACCACATCCTGTGCTGGCCGCTGGGGGCACCAG  
GTGAGGTGGTGGCTGTGCCCTGTCCGACT  
ACATTTATGACTTCAATCACAAAGGCCATGCCTACCGACGCTGTGACCGCAA  
TGGCAGCTGGGAGCTGGTGCCTGGGCAC  
AACAGGACGTGGGCCAACTACAGCGAGTGTGTCAAATTTCTCACCAATGAGA  
CTCGTGAACGGGAGGTGTTTGACCGCCT  
GGGCATGATTTACACCGTGGGCTACTCCGTGTCCCTGGCGTCCCTCACCGT  
AGCTGTGCTCATCCTGGCCTACTTTAGGC  
GGCTGCACTGCACGCGCAACTACATCCACATGCACCTGTTCCCTGTCCTTCAT  
GCTGCGCGCCGTGAGCATCTTCGTCAAG  
GACGCTGTGCTCTACTCTGGCGCCACGCTTGATGAGGCTGAGCGCCTCACC  
GAGGAGGAGCTGCGCGCCATCGCCCAGGC  
GCCCCCGCCGCCTGCCACCGCCGCTGCCGGCTACGCGGGCTGCAGGGTG  
GCTGTGACCTTCTTCCTTTACTTCCTGGCCA  
CCAACTACTACTGGATTCTGGTGGAGGGGCTGTACCTGCACAGCCTCATCTT  
CATGGCCTTCTTCTCAGAGAAGAAGTAC  
CTGTGGGGCTTCACAGTCTTCGGCTGGGGTCTGCCCGCTGTCTTCGTGGCT  
GTGTGGGTCAGTGTGAGAGCTACCCTGGC  
CAACACCGGGTGCTGGGACTTGAGCTCCGGGAACAAAAAGTGGATCATCCA  
GGTGCCCATCCTGGCCTCCATTGTGCTCA  
ACTTCATCCTCTTCATCAATATCGTCCGGGTGCTCGCCACCAAGCTGCGGGA  
GACCAACGCCGGCCGGTGTGACACACGG  
CAGCAGTACCGGAAGCTGCTCAAATCCACGCTGGTGCTCATGCCCCTCTTTG  
CGGTCCACTACATTGTCTTCATGGCCAC  
ACCATACACCGAGGTCTCAGGGACGCTCTGGCAAGTCCAGATGCACTATGA  
GATGCTCTTCAACTCCTTCAGGGATTTT  
TTGTCGCAATCATATACTGTTTCTGCAATGGCGAGGTACAAGCTGAGATCAA  
GAAATCTTGGAGCCGCTGGACACTGGCA  
CTGGACTTCAAGCGAAAGGCACGCAGCGGGAGCAGCAGCTATAGCTACGGC  
CCCATGGTGTCCACACAAGTGTGACCAA  
TGTCGGCCCCCGTGTGGGACTCGGCCTGCCCCTCAGCCCCCGCCTACTGC  
CCACTGCCACCACCAACGGCCACCCTCAGC

TGCCTGGCCATGCCAAGCCAGGGACCCCAGCCCTGGAGACCCTCGAGACC  
 ACACCACCTGCCATGGCTGCTCCCAAGGAC  
 GATGGGTTCTCAACGGCTCCTGCTCAGGCCTGGACGAGGAGGCCTCTGGG  
 CCTGAGCGGCCACCTGCCCTGCTACAGGA  
 AGAGTGGGAGACAGTCATGTGATGA

**SEQ ID NO: 4: wildtype Human PTH/PTHrP receptor amino acid sequence:**

MGTARIAPGLALLCCPVLSSAYALVDADDVMTKEEQIFLLHRAQAQCEKRLKEV  
 LQRPASIMESDKGWTSASTSGKPRK  
 DKASGKLYPESEEDKEAPTGSRYRGRPCLEWDHILCWPLGAPGEVVAVPCPD  
 YIYDFNHKGHAYRRCDRNGSWELVPGH  
 NRTWANYSECVKFLTNETREREVEFDRLGMIYTVGYSVSLASLTAVLILAYFRRL  
 HCTRNYIHMHLFLSFMLRAVSIFVK  
 DAVLYSGATLDEAERLTEEELRAIAQAPPPATAAAGYAGCRVAVTFFLYFLATN  
 YYWILVEGLYLHSLIFMAFFSEKKY  
 LWGFTVFGWGLPAVFVAVWVSVRATLANTGCWDLSSGNKKWIIQVPILASIVLN  
 FILFINIVRVLATKLRETNAGRCDDR  
 QQYRKLLKSTLVLMPLFGVHYIVFMATPYTEVSGTLWQVQMHYEMLFNSFQGFF  
 VAIYCFCNGEVQAEIKKSWSRWTLA  
 LDFKRKARSGSSSYSGPMVSHTSVTNVGPRVGLGLPLSPRLLPTATTNGHPQL  
 PGHAKPGTPALETLETPPAMAAPKD  
 DGFLNGSCSGLDEEASGPERPPALLQEEWETVM

**SEQ ID NO: 5: wildtype A2A adenosine receptor (human) cDNA: Genbank:**  
***M97370 (corresponding to mouse: XM 125720)***

ATGCCCATCATGGGCTCCTCGGTGTACATCACGGTGGAGCTGGCCATTGCT  
 GTGCTGGCCATCCTGGGCAATGTGCTGGTGTGCTGGGCCGTGTGGCTCAAC  
 AGCAACCTGCAGAACGTCACTCACTTTGTGGTGTCACTGGCGGGCGGCC  
 GACATCGCAGTGGGTGTGCTCGCCATCCCCTTTGCCATCACCATCAGCACC  
 GGGTTCTGCGCTGCCTGCCACGGCTGCCTCTTCATTGCCTGCTTCGTCCTG

GTCTCACGCAGAGCTCCATCTTCAGTCTCCTGGCCATCGCCATTGACCGCT  
ACATTGCCATCCGCATCCCGCTCCGGTACAATGGCTTGGTGACCGGCACGA  
GGGCTAAGGGGCATCATTGCCATCTGCTGGGTGCTGTCGTTTGCCATCGGCC  
TGA TCCCATGCTAGGTTGGAACA ACTGCGGTCAGCCAAAGGAGGGCAAGA  
ACCACTCCCAGGGCTGCGGGGAGGGCCAAAGTGGCCTGTCTCTTTGAGGATG  
TGGTCCCCATGAACTACATGGTGTACTTCAACTTCTTTCCTGTGTGCTGGTGC  
CCCTGCTGCTCATGCTGGGTGTCTATTTGCGGATCTTCCTGGCGGGCGCGAC  
GACAGCTGAAGCAGATGGAGAGCCAGCCTCTGCCGGGGGAGCGGGGCACGG  
TCCACACTGCAGAAGGAGGTCCATGCTGCCAAGTCACTGGCCATCATTGTG  
GGGCTCTTTGCCCTCTGCTGGCTGCCCCTACACATCATCAACTGCTTCACTT  
TCTTCTGCCCCGACTGCAGCCACGCCCCCTCTCTGGCTCATGTACCTGGCCAT  
CGTCCTCTCCCACACCAATTGCGTTGTGAATCCCTTCATCTACGCCTACCGT  
ATCCGCGAGTTCCGCCAGACCTTCGCAAGATCATTGCGAGCCACGTCCTG  
AGGCAGCAAGAACCTTTCAAGGCAGCTGGCACCAGTGCCCGGGTCTTGGA  
GCTCATGGCAGTGACGGAGAGCAGGTGAGCCTCCGTCTCAACGGCCACCCG  
CCAGGAGTGTGGGCCAACGGCAGTGCTCCCCACCCTGAGCGGAGGCCCAA  
TGGCTATGCCCTGGGGCTGGTGAGTGGAGGGAGTGCCCAAGAGTCCCAGG  
GGAACACGGGCCTCCCAGACGTGGAGCTCCTTAGCCATGAGCTCAAGGGAG  
TGTGCCCAGAGCCCCCTGGCCTAGATGACCCCCTGGCCCAGGATGGAGCA  
GGAGTGTCTGA

**SEQ ID NO: 6: wildtype A2A adenosine receptor (human) amino acid sequence**

MPIMGSSVYITVELAI AVLAILGNVLCWAVWLNSNLQNV TNYFVSLAAADIAVGV  
LAIPFAITISTGFCAACHGCLFIACFVLVLTQSSIFSLLAIAIDRYIAIRIPLRYNGLVT  
GTRAKGIIAICWVLSFAIGLTPMLGWNNCGQPKEGKNHSQGC GEGQVACLFEDV  
VPMNYMVYFNFFACVLVPLLLMLGVYLRIFLAARRQLKQMESQPLPGERARSTL  
QKEVHAAKSLAIIVGLFALCWLPLHIINCFTFFCPDCSHAPLWLMYLAIVLSHTNSV  
VNPFIYAYRIREFRQTFRKIIRSHVLRQQEPFKAAGTSARVLA AHGSDGEQVSLR  
LNGHPPGVWANGSAPHPERRPNGYALGLVSGGSAQESQGNTGLPDVELLSHE  
LKG

The following fluorophore sequences have been employed in the appended examples:

**SEQ ID NO: 7: eCFP (enhanced CFP) cDNA sequence (*Clontech*)**

ATGGTGAGCAAGGGGCGAGGAGCTGTTACCGGGGTGGTGCCCATCCTGGT  
CGAGCTGGACGGCGACGTAAACGGCCACAA  
GTTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGA  
CCCTGAAGTTCATCTGCACCACCGGCAAGC  
TGCCCGTGCCCTGGCCCAACCCTCGTGACCACCCTGACCTGGGGCGTGCACT  
GCTTCAGCCGCTACCCCGACCACATGAAG  
CAGCACGACTTCTTCAAGTCCGCCATGCCCCAAGGCTACGTCCAGGAGCGC  
ACCATCTTCTTCAAGGACGACGGCAACTA  
CAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCA  
TCGAGCTGAAGGGCATCGACTTCAAGGAGG  
ACGGCAACATCCTGGGGCACAAGCTGGAGTACAACCTACATCAGCCACAACG  
TCTATATCACCGCCGACAAGCAGAAGAAC  
GGCATCAAGGCCAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTG  
CAGCTCGCCGAACCACTACCAGCAGAACAC  
CCCCATCGGGCGACGGCCCCGTGCTGCTGCCCCGACAACCACTACCTGAGCAC  
CCAGTCCGCCCTGAGCAAAGACCCCAACG  
AGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGACCGCCGCGGGGATCA  
CTCTCGGCATGGACGAGCTGTACAAGTAA

**SEQ ID NO: 8: eCFP (enhanced CFP) amino acid sequence:**

MVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTGKLP  
VPWPTLVTTLTWGVQCFSRYPDHMK  
QHDFFKSAMPEGYVQERTIFFKDDGNYKTRAIEVKFEGDTLVNRIELKGIDFKEDG  
NILGHKLEYNYSIHNVYITADKQKN

GIKANFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSALSKDPNEK  
RDHMLLEFVTAAGITLGMDELYK

**SEQ ID NO: 9: YFP cDNA sequence: (Clontech)**

ATGGTGAGCAAGGGCGAGGAGCTGTTCACCGGGGTGGTGCCCATCCTGGT  
CGAGCTGGACGGCGACGTAAACGGCCACAA  
GTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGA  
CCCTGAAGTTCATCTGCACCACCGGCAAGC  
TGCCCGTGCCCTGGCCCAACCCTCGTGACCACCTTCGGCTACGGCCTGCAGT  
GCTTCGCCCCGCTACCCCGACCACATGAAG  
CAGCAGGACTTCTTCAAGTCCGCCATGCCGAAGGCTACGTCCAGGAGCGC  
ACCATCTTCTTCAAGGACGACGGCAACTA  
CAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCA  
TCGAGCTGAAGGGCATCGACTTCAAGGAGG  
ACGGCAACATCCTGGGGCACAAGCTGGAGTACAACACTACAACAGCCACAACG  
TCTATATCATGGCCGACAAGCAGAAGAAC  
GGCATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTG  
CAGCTCGCCGACCACTACCAGCAGAACAC  
CCCCATCGGCGACGGCCCCGTGCTGCTGCCCGACAACCACTACCTGAGCTA  
CCAGTCCGCCCTGAGCAAAGACCCCAACG  
AGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGACCGCCGCCGGGATCA  
CTCTCGGCATGGACGAGCTGTACAAGTAA

**SEQ ID NO: 10: YFP amino acid sequence:**

MVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTGKLP  
VPWPTLVTTFGYGLQCFARYPDHMK  
QHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDG  
NILGHKLEYNYNshNVYIMADKQKN  
GIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHLSYQSALSKDPNEK  
RDHMLLEFVTAAGITLGMDELYK

In the following receptor-cam ("chameleon") sequences the CFP and YFP sequences are identified by bold printing:

SEQ ID NO: 11: alpha2a adrenergic receptor-cam cDNA sequence:

ATGGGCTACCCATACGACGTCCCAGACTACGCCAGCATGGGCTCACTGCAG  
 CCGGATGCCGGCAACAGCAGCTGGAACGG  
 GACCGAAGCGCCCGGAGGCGGCACCCGAGCCACCCCTTACTCCCTGCAGG  
 TGACACTGACGCTGGTTTGCCTGGCTGGCC  
 TGCTCATGCTGTTACAGTATTTGGCAACGTGCTGGTTATTATCGCGGTGTTT  
 ACCAGTCGCGCGCTCAAAGCTCCCCAA  
 AACCTCTTCCTGGTGTCCCTGGCCTCAGCGGACATCCTGGTGGCCACGCTG  
 GTCATTCCCTTTTCTTTGGCCAACGAGGT  
 TATGGGTTACTGGTACTTTGGTAAGGTGTGGTGTGAGATCTATTTGGCTCTC  
 GACGTGCTCTTTTGCACGTCGTCCATAG  
 TGCACCTGTGCGCCATCAGCCTTGACCGCTACTGGTCCATCACGCAGGCCA  
 TCGAGTACAACCTGAAGCGCACGCCGCGT  
 CGCATCAAGGCCATCATTGTCACCGTGTGGGTGTCATCTCGGCTGTCATCTCCT  
 TCCCGCCACTCATCTCCATAGAGAAGAA  
 GGGCGCTGGCGGGCGGGCAGCAGCCGGCCGAGCCAAGCTGCAAGATCAACG  
 ACCAGAAGTGGTATGTCATCTCCTCGTCCA  
 TCGGTTCCCTTCTTCGCGCCTTGCCTCATCATGATCCTGGTCTACGTGCGTATT  
 TACCAGATCGCCAAGCGTCGCACCCGC  
GTGCCTCCCAGCCGCGGGGTCCGGACGCCATGGTGAGCAAGGGCGAGG  
AGCTGTTACACGGGGTGGTGCCCATCCTGGT  
CGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAG  
GGCGAGGGCGATGCCACCTACGGCAAGCTGA  
CCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCGTGCCCTGGCCCACC  
CTCGTGACCACCTTCGGCTACGGCCTGCAG  
TGCTTCGCCCCGCTACCCCGACCACATGAAGCAGCACGACTTCTTCAAGTCC  
GCCATGCCCCGAAGGCTACGTCCAGGAGCG

CACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGA  
AGTTCGAGGGCGACACCCTGGTGAACCGCA  
TCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCAC  
AAGCTGGAGTACAACACTACAACAGCCACAAC  
GTCTATATCATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACTTCAA  
GATCCGCCACAACATCGAGGACGGCAGCGT  
GCAGCTCGCCGACCACTACCAGCAGAACACCCCATCGGCGACGGCCCC  
GTGCTGCTGCCCGACAACCACTACCTGAGCT  
ACCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTC  
CTGCTGGAGTTCGTGACCGCCGCGGGGATC  
ACTCTCGGCATGGACGAGCTGTACAAGCGCTGGCGCGGGAGGCAAAACCG  
GGAGAAACGCTTCACGTTCTGTGCTGGCGGT  
GGTGATCGGCGTGTTCTGTGGTGTGTTGGTTTCCGTTCTTTTTACCTACACG  
CTCATAGCGGTGCGCTGCCCGGTGCCCA  
GCCAGCTCTTCAACTTCTTCTTCTGGTTCGGCTACTGCAACAGCTCGCTGAA  
CCCTGTTATCTACACCATCTTCAACCAC  
GACTTCCGACGCGCCTTCAAGAAGATCCTCTGCCGTGGGGACAGAAAACGC  
ATCGTGATGGTGAGCAAGGGCGAGGAGCT  
GTTACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACGTAAACG  
GCCACAAGTTCAGCGTGTCCGGCGAGGGCG  
AGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACC  
GGCAAGCTGCCCGTGCCCTGGCCACCCCTC  
GTGACCACCCTGACCTGGGGCGTGCAAGTGCTTCAGCCGCTACCCCGACCA  
CATGAAGCAGCAGACTTCTTCAAGTCCGC  
CATGCCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACG  
GCAACTACAAGACCCGCGCCGAGGTGAAGT  
TCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTC  
AAGGAGGACGGCAACATCCTGGGGCACAAG  
CTGGAGTACAACACTACATCAGCCACAACGTCTATATCACCGCCGACAAGCA  
GAAGAACGGCATCAAGGCCAACTTCAAGAT  
CCGCCACAACATCGAGGACGGCAGCGTGCAAGCTCGCCGACCACTACCAGC  
AGAACACCCCATCGGCGACGGCCCCGTGC

TGCTGCCCGACAACCACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGAC  
CCCAACGAGAAGCGCGATCACATGGTCCTG  
CTGGAGTTCGTGACCGCCGCCGGGATCACTCTCGGCATGGACGAGCTGTA  
CAAGTAA

SEQ ID NO: 12: Alpha2a adrenergic receptor-cam "chameleon" amino acid sequence:

MGYPYDVPDYASMGSLQPDAGNSSWNGTEAPGGGTRATPYSLQVTLTLVCLA  
 GLLMLFTVFGNVLVIIAVFTSRALKAPQ  
 NLFLVSLASADILVATLVIPFSLANEVMGYWYFGKVCCEIYLALDVLFTSSIVHLC  
 AISLDRYWSITQAIEYNLKRTPR  
 RIKAIIVTVWVISAVISFPPLISIEKKGAGGGQQPAEPSCKINDQKWYVISSSIGSFF  
 APCLIMILVYVRIYQIAKRRT  
VPPSRRGPDAMVSKGEELFTGVVPILVELDGDVNGHKFSVSgegeDATYGKL  
TLKFICTTGKLPVPWPTLVTTFGYGLQ  
CFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVN  
RIELKGIDFKEDGNILGHKLEYNNSHN  
VYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSY  
QSALSKDPNEKRDHMLLEFVTAAGI  
TLGMDELYKRWGRGRQREKRFTFVLAVVIGVFVVCWFPFFFTYTLIAVGCPVPS  
 QLFNFFFWFGYCNSSLNPVIYTIFNH  
 DFRRAFKKILCRGDRKRIVMVSKGEELFTGVVPILVELDGDVNGHKFSVSgege  
GDATYGKLTTLKFICTTGKLPVPWPTL  
VTTLTWGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEV  
KFEGDTLVNRIELKGIDFKEDGNILGHK  
LEYNYISHNVYITADKQKNGIKANFKIRHNIEDGSVQLADHYQQNTPIGDGPVLL  
PDNHYSTQSALSKDPNEKRDHML  
LEFVTAAGITLGMDELYK

SEQ ID NO: 13: PTH receptor-cam "chameleon" 8 cDNA sequence:



ATGGGGACCGCCCGGATCGCACCCGGCCTGGCGCTCCTGCTCTGCTGCCC  
CGTGCTCAGCTCCGCGTACGCGCTGGTGGA  
TGCAGATGACGTCATGACTAAAGAGGAACAGATCTTCCTGCTGCACCGTGCT  
CAGGCCCAAGTGCAGAAAACGGCTCAAGG  
AGGTCCTGCAGAGGCCAGCCAGCATAATGGAATCAGACAAGGGATGGACAT  
CTGCGTCCACATCAGGGAAGCCCAGGAAA  
GATAAGGCATCTGGGAAGCTCTACCCTGAGTCTGAGGAGGACAAGGAGGCA  
CCCACTGGCAGCAGGTACCGAGGGCGCCC  
CTGTCTGCCGGAATGGGACCACATCCTGTGCTGGCCGCTGGGGGCACCAG  
GTGAGGTGGTGGCTGTGCCCTGTCCGGACT  
ACATTTATGACTTCAATCACAAAGGCCATGCCTACCGACGCTGTGACCGCAA  
TGGCAGCTGGGAGCTGGTGCCTGGGCAC  
AACAGGACGTGGGCCAACTACAGCGAGTGTGTCAAATTTCTACCAATGAGA  
CTCGTGAACGGGAGGTGTTTGACCGCCT  
GGGCATGATTTACACCGTGGGCTACTCCGTGTCCCTGGCGTCCCTCACCGT  
AGCTGTGCTCATCCTGGCCTACTTTAGGC  
GGCTGCACTGCACGCGCAACTACATCCACATGCACCTGTTCTGTCTTCAT  
GCTGCGCGCCGTGAGCATCTTCGTCAAG  
GACGCTGTGCTCTACTCTGGCGCCACGCTTGATGAGGCTGAGCGCCTCACC  
GAGGAGGAGCTGCGCGCCATCGCCCAGGC  
GCCCCGCGCGCTGCCACCGCCGCTGCCGGCTACGCGGGCTGCAGGGTG  
GCTGTGACCTTCTTCCTTTACTTCCTGGCCA  
CCAATACTACTGGATTCTGGTGGAGGGGCTGTACCTGCACAGCCTCATCTT  
CATGGCCTTCTTCTCAGAGAAGAAGTAC  
CTGTGGGGCTTCACAGTCTTCGGCTGGGGTCTGCCCCGCTGTCTTCGTGGCT  
GTGTGGGTGAGTGTGAGAGCTACCCTGGC  
CAACACCGGGTGCTGGGACTTGAGCTCCGGGAACAAAAAGTGGATCATCCA  
GGTGCCCATCCTGGCCTCCATTGTGCTCA  
ACTTCATCCTCTTCATCAATATCGTCCGGGTGCTCGCCACCAAGCTGCGGGA  
GACCAACGCCGGCATGGTGAGCAAGGGC  
GAGGAGCTGTTACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGA  
CGTAAACGGCCACAAGTTCAGCGTGTCCGG

CGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCT  
GCACCACCGGCAAGCTGCCCGTGCCCTGGC  
CCACCCTCGTGACCACCTGACCTGGGGCGTGACGTGCTTCAGCCGCTAC  
CCCGACCACATGAAGCAGCAGCACTTCTTC  
AAGTCCGCCATGCCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAA  
GGACGACGGCAACTACAAGACCCGCGCCGA  
GGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCA  
TCGACTTCAAGGAGGACGGCAACATCCTGG  
GGCACAAGCTGGAGTACAACCTACATCAGCCACAACGTCTATATCACCGCC  
GACAAGCAGAAGAACGGCATCAAGGCCAAC  
TTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGACGCTCGCCGACCA  
CTACCAGCAGAACACCCCATCGGCGACGG  
CCCCGTGCTGCTGCCCGACAACCACTACCTGAGCACCCAGTCCGCCCTGA  
GCAAAGACCCCAACGAGAAGCGCGATCACA  
TGGTCCTGCTGGAGTTCGTGACCGCCGCGGGATCACTCTCGGCATGGAC  
GAGCTGTACAAGCGGTGTGACACACGGCAG  
CAGTACCGGAAGCTGCTCAAATCCACGCTGGTGCTCATGCCCCTCTTTGGC  
GTCCACTACATTGTCTTCATGGCCACACC  
ATACACCGAGGTCTCAGGGACGCTCTGGCAAGTCCAGATGCACTATGAGAT  
GCTCTTCAACTCCTTCCAGGGATTTTTTG  
TCGCAATCATATACTGTTTCTGCAATGGCGAGGTACAAGCTGAGATCAAGAA  
ATCTTGGAGCCGCTGGACACTGGCACTG  
GACTTCAAGCGAAAGGCACGCAGCGGGAGCAGCAGCTATAGCTACGGCATG  
GTGAGCAAGGGCGAGGAGCTGTTACCGG  
GGTGGTGCCCATCCTGGTTCGAGCTGGACGGCGACGTAAACGGCCACAAGT  
TCAGCGTGTCGGCGAGGGCGAGGGCGATG  
CCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGCAAGCTG  
CCCGTGCCCTGGCCACCCCTCGTGACCACC  
TTGGGCTACGGCCTGCAGTGCTTCGCCCCGCTACCCCGACCACATGAAGCA  
GCACGACTTCTTCAAGTCCGCCATGCCCGA  
AGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACA  
AGACCCGCGCCGAGGTGAAGTTCGAGGGCG

ACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGAC  
GGCAACATCCTGGGGCACAAGCTGGAGTAC  
AACTACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAACGG  
CATCAAGGTGAACTTCAAGATCCGCCACAA  
CATCGAGGACGGCAGCGTGCAGCTCGCCGACCACTACCAGCAGAACACCC  
CCATCGGGCGACGGGCCCCGTGCTGCTGCCCG  
ACAACCACTACCTGAGCTACCAGTCCGCCCTGAGCAAAGACCCCAACGAG  
AAGCGCGATCACATGGTCCTGCTGGAGTTC  
GTGACCGCCGCCGGGATCACTCTCGGCATGGACGAGCTGTACAAGTAA

SEQ ID NO: 14: PTH receptor-cam8 "chameleon" amino acid sequence:

MGTARIAPGLALLCCPVLSSAYALVDADDVMTKEEQIFLLHRAQAQCEKRLKEV  
 LQRPASIMESDKGWTSASTSGKPRK  
 DKASGKLYPESEEDKEAPTGSRYRGRPCLEWDHILCWPLGAPGEVVAVPCPD  
 YIYDFNHKGHAYRRCDRNGSWELVPGH  
 NRTWANYSECVKFLTNETREREVFDRLGMIYTVGYSVSLASLTVAVLILAYFRRL  
 HCTRNYIHMHLFLSFMLRAVSIFVK  
 DAVLYSGATLDEAERLTEEELRAIAQAPPPATAAAGYAGCRVAVTFFLYFLATN  
 YYWILVEGLYLHSLIFMAFFSEKKY  
 LWGFTVFGWGLPAVFVAVVWSVRATLANTGCWDLSSGNKKWIIQVPILASIVLN  
 FILFINIVRVLATKLRETNAGMVSKG  
EELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICTTGKLPVPWP  
 TLVTTLTWGVQCFSRYPDHMKQHDF  
KSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILG  
 HKLEYNVISHNVYITADKQKNGIKAN  
FKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSQSALSKDPNEKRDH  
 MVLLEFVTAAGITLGMDELYKRC| Q | Y | R | K | L | L | K | S | T | L | V | L | M | P | L | F | G | V | H | I | V | F | M | A | T | P | Y | T | E | V | S | G | T | L | W | Q | V | Q | M | H | Y | E | M | L | F | N | S | F | Q | G | F | F | V |
 A | I | I | Y | C | F | C | N | G | E | V | Q | A | E | I | K | K | S | W | S | R | W | T | L | A | L | D | F | K | R | K | A | R | S | G | S | S | S | Y | S | Y | G | M | V | S | K | G | E | E | L | F | T | G | V | P | I | L | V | E | L | D | G | D | V | N | G | H | K | F | S | V | S | G | E | G | E | G | D | A | T | Y | G | K | L | T | L | K | F | I | C | T | T | G | K | L | P | V | P | W | P | T | L | V | T |

FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY  
NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALS KDPNEKRDHMLLEF  
VTAAGITLGMDELYK

SEQ ID NO: 15: A2A-CFP14/10-YFP-C33 cDNA sequence; A2A "chameleon"

ATGCCCATCATGGGCTCCTCGGTGTACATCACGGTGGAGCTGGCCATTG  
CTGTGCTGGCCATCCTGGGCAATGTGCTGGTGTGCTGGGCCGTGTGGCTC  
AACAGCAACCTGCAGAACGTCACCAACTACTTTGTGGTGTCACTGGCGGC  
GGCCGACATCGCAGTGGGTGTGCTCGCCATCCCCTTTGCCATCACCATCA  
GCACCGGGTTCTGCGCTGCCTGCCACGGCTGCCTCTTCATTGCCTGCTTCG  
TCCTGGTCC TCACGCAGAG CTCCATCTTC AGTCTCCTGG CCATCGCCAT  
TGACCGCTAC ATTGCCATCCGCATCCCGCT CCGGTACAAT GGCTTGGTGA  
CCGGCACGAG GGCTAAGGGC ATCATTGCCATCTGCTGGGTGCTGTGCTTT  
GCCATCGGCC TGA CTCCCATGCTAGGTTGGAACA ACTGCGGTCAGCCAAA  
GGAGGGGCAAGAACCACTCCCAGGGGCTGCGGGGAGGGCCAAGTGGCCTGTC  
TCTTTGAGGA TGTGGTCCCC ATGAACTACATGGTGTACTTCAACTTCTTT  
GCCTGTGTGCTGGTGCCCT GCTGCTCATG CTGGGTGTCT ATTTGCGGAT  
CTTCCTGGCGGCGCGACGACAGCTGAAGCAGATGGAGGTGAGCAAGGGCG  
AGGAGCTGTTACCGGGGGTGGTGCCCATCCTGGTTCGAGCTGGACGGCGAC  
GTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGGCGAGGGCGATGCCA  
CCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCAACCGGCAAGCTGCCC  
GTGCCCTGGCCCACCCTCGTGACCACCCTGACCTGGGGCGTGCAAGTGCTT  
CAGCCGCTACCCCGACCATGAAGCAGCAGACTTCTTCAAGTCCGCCA  
TGCCCCAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGC  
AACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGGCGACACCCTGGTGAA  
CCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGG  
GGCACAAGCTGGAGTACA ACTACATCAGCCACAACGTCTATATCACCGCC  
GACAAGCAGAAGAACGGCATCAAGGCCAACTTCAAGATCCGCCACAACAT  
CGAGGACGGCAGCGTGCAAGCTCGCCGACCACTACCAGCAGAACACCCCC

ATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACCACTACCTGAGCACCCA  
GTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCCTGC  
TGGAGTTCGTGACCGCCGCGGGGATCACTCTCGGCATGGACGAGCTGTAC  
AAG CTTCAGA AGGAGGTCCA TGCTGCCAAG TCACTGGCCA  
 TCATTGTGGG GCTCTTTGCC CTCTGCTGGC TGCCCCTACA CATCATCAAC  
 TGCTTCACTT TCTTCTGCCC CGACTGCAGC CACGCCCCTC TCTGGCTCAT  
 GTACCTGGCC ATCGTCCTCT CCCACACCAA TTCGGTTGTG AATCCCTTCA  
 TCTACGCCTA  
 CCGTATCCGCGAGTTCCGCCAGACCTTCCGCAAGATCATTGCGAGCCACGT  
 CCTGAGGCAGCAAGAACCTTTCAAGGCAG CTGGCACCAGTGCCCCGGGTC  
GTGAGCAAGGGCGAGGAGCTGTTACCGGGGTGGTGCCCATCCTGGT  
CGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAG  
GGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCAC  
CACCGGCAAGCTGCCCGTGCCCTGGCCACCCCTCGTGACCACCTTCGGCT  
ACGGCCTGCAG  
TGCTTCGCCCCGCTACCCCGACCACATGAAGCAGCACGACTTCTTCAAGTCC  
GCCATGCCCGAAGGCTACGTCCAGGAGCG  
CACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGA  
AGTTCGAGGGCGACACCCTGGTGAACCGCA  
TCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCAC  
AAGCTGGAGTACAACACTACAACAGCCACAAC  
GTCTATATCATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACTTCAA  
GATCCGCCACAACATCGAGGACGGCAGCGT  
GCAGCTCGCCGACCACTACCAGCAGAACACCCCATCGGGCGACGGCCCC  
GTGCTGCTGCCCCGACAACCACTACCTGAGCT  
ACCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTC  
CTGCTGGAGTTCGTGACCGCCGCGGGGATC  
 ACTCTCGGCATGGACGAGCTGTACAAG

SEQ ID NO: 16: A2A-CFP14/10-YFP-C33 amino acid sequence; "A2A  
 chameleon"

MPIMGSSVYITVELAIAVLAILGNVLVCWAVWLNSNLQNVNTNYFVSLAAADIAVGV  
 LAIPFAITISTGFCAACHGCLFIACFVLVLTQSSIFSLLAIAIDRYIAIRIPLRYNGLVT  
 GTRAKGIIAICWVLSFAIGLTPMLGWNNCGQPKEGKNHSQGC GEGQVACLFEDV  
 VPMNYMVYFNFFACVLVPLLLMLGVYLRIFLAARRQLKQMEVSKGEELFTGVVPI  
LVELDGDVNGHKFSVS GEGEGDATYGKLT LKFICTTGKLPVPWPTLVTTLTWG  
VQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTL  
VNRIELKGIDFKEDGNILGHKLEYN YISHNVYITADKQKNGIKANFKIRHNIEDGS  
VQLADHYQQNTPIGDGPVLLPDNHYSTQSALSKDPNEKRDHMLLEFVTAAG  
 ITLGMDELYKLQKEVHAAKSLAIVGLFALCWLPLHIINCFTFFCPDCSHAPLWLM  
 YLAIVLSHTNSVNPFIYAYRIREFRQTFRKIIRSHVLRQQEPFKAAGTSARVMVS  
KGEELFTGVVPILVELDGDVNGHKFSVS GEGEGDATYGKLT LKFICTTGKLPVP  
WPTLVTTFGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKT  
RAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYN YNSHNVYIMADKQKNGIK  
VNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYSYQSALSKDPNEKRD  
HMLLEFVTAAGITLGMDE

Further chameleon-constructs; in particular, PTH-constructs are defined by appended SEQ ID NOS: 29 to 37. A control-construct as employed in accordance with this invention is appended in SEQ ID NO: 38.

Loops and C-termini to be modified for chameleon-constructs of the invention are exemplified in the following SEQ ID NOS: 17 to 28.

**SEQ ID NO: 17: Mouse  $\alpha$ 2a adrenergic receptor third intracellular loop cDNA sequence:**

GTGCGTATTTACCAGATCGCCAAGCGTCGCACCCGCGTGCCTCCCAGCCGC  
 CGGGGTCCGGACGCCTGTTCCGCGCCGCC  
 GGGGGGCGCCGATCGCAGGCCCAACGGGCTGGGCCCGGAGCGCGGCGCG  
 GGTCCCACGGGCGCTGAGGCGGAGCCGCTGC  
 CCACCCAGCTTAACGGTGCCCCGGGGGAGCCCGCGCCCGCGGGCCCCG  
 CGATGGGGATGCGCTGGACCTAGAGGAGAGT

TCGTCGTCCGAGCACGCCGAGCGGCCCCCGGGGCCCCGCAGACCCGACCG  
 CGGCCCCCGAGCCAAGGGCAAGACCCGGGC  
 GAGTCAGGTGAAGCCGGGGGACAGTCTGCCGCGGCGCGGGCCCCGGGGCC  
 GCGGGGCCGGGGGCTTCGGGGTCCGGGCACG  
 GAGAGGAGCGCGGCGGGGGCGCCAAAGCGTCGCGCTGGCGCGGGAGGCA  
 AAACCGGGAGAAACGCTTCACGTTCGTG

**SEQ ID NO: 18: Mouse alpha2a adrenergic receptor third Intracellular loop amino acid sequence:**

VRIYQIAKRRTRVPPSRRGPDACSAPPGGADRRPNGLGPERGAGPTGAEEPL  
 PTQLNGAPGEPAPAGPRDGDALDLEES  
 SSSEHAERPPGPRRPDRGPRAKGKTRASQVKPGDSLPRRGPGAAGPGASGSG  
 HGEERGGGAKASRWRGRQNREKRFTFV

**SEQ ID NO: 19: Mouse alpha2a adrenergic receptor carboxy terminus tail cDNA sequence:**

CACGACTTCCGACGCGCCTTCAAGAAGATCCTCTGCCGTGGGGACAGAAAA  
 CGCATCGTGTGA

**SEQ ID NO: 20: Mouse alpha2a adrenergic receptor carboxy terminus tail amino acid sequence:**

HDFRRAFKKILCRGDRKRIV

**SEQ ID NO. 21: human A2A adenosine receptor 3<sup>rd</sup>.loop sequence cDNA:**

GGATCTTCCTGGCGGCGCGACGACAGCTGAAGCAGATGGAGAGCCAGCCT  
 CTGCCGGGGGAGCGGGCACGGTCCACACTGCAGAAGGAGGTCCATGCTGC  
 CAAGTCA

**SEQ ID NO: 22: human A2A adenosine receptor 3<sup>rd</sup>.loop amino acid sequence :**

RIFLAARRQLKQMESQPLPGERARSTLQKEVHAAKS

**SEQ ID NO: 23: human A2A adenosine receptor c-terminus cDNA sequence:**

CGTATCCGCGAGTTCCGCCAGACCTTCCGCAAGATCATTTCGCAGCCACGTC  
CTGAGGCAGCAAGAACCTTTCAAGGCAGCTGGCACCAAGTGCCCCGGGTCTTG  
GCAGCTCATGGCAGTGACGGAGAGCAGGTCAGCCTCCGTCTCAACGGCCAC  
CCGCCAGGAGTGTGGGCCAACGGCAGTGCTCCCCACCCTGAGCGGAGGCC  
CAATGGCTATGCCCTGGGGCTGGTGAGTGGAGGGAGTGCCCAAGAGTCCCA  
GGGGAACACGGGCCTCCCAGACGTGGAGCTCCTTAGCCATGAGCTCAAGG  
GAGTGTGCCCAGAGCCCCCTGGCCTAGATGACCCCCTGGCCCAGGATGGA  
GCAGGAGTGTCTGA

**SEQ ID NO: 24: human A2A adenosine receptor c-terminus amino acid sequence:**

RIREFRQTFRKIIRSHVLRQQEPFKAAGTSARVLAHGS DGEQVSLRLNGHPPG  
VWANGSAPHPERRPNGYALGLVSGGSAQESQGNTGLPDVELLSHELKG

**SEQ ID NO: 25: human PTH/PTHrP receptor third intracellular loop cDNA sequence:**

ACCAAGCTGCGGGAGACCAACGCCGGCCGGTGTGACACACGGCAGCAGTA  
CCGGAAG

**SEQ ID NO: 26: human PTH/PTHrP receptor third intracellular loop amino acid sequence:**

TKLRETNAGRC DTRQQYRK



**SEQ ID NO: 27: human PTH/PTHrP receptor carboxy terminus cDNA sequence:**

GAGGTACAAGCTGAGATCAAGAAATCTTGGAGCCGCTGGACACTGGCACTG  
 GACTTCAAGCGAAAGGCACGCAGCGGGAG  
 CAGCAGCTATAGCTACGGCCCCATGGTGTCCCACACAAGTGTGACCAATGTC  
 GGCCCCCGTGTGGGACTCGGCCTGCCCC  
 TCAGCCCCCGCCTACTGCCCACTGCCACCACCAACGGCCACCCTCAGCTGC  
 CTGGCCATGCCAAGCCAGGGACCCCAGCC  
 CTGGAGACCCTCGAGACCACACCACCTGCCATGGCTGCTCCCAAGGACGAT  
 GGGTTCCTCAACGGCTCCTGCTCAGGCCT  
 GGACGAGGAGGCCTCTGGGCCTGAGCGGCCACCTGCCCTGCTACAGGAAG  
 AGTGGGAGACAGTCATGTGATGA

**SEQ ID NO: 28: human PTH/PTHrP receptor carboxy terminus amino acid sequence:**

EVQAEIKKSWSRWTLALDFKRKARSGSSSYSGPMVSHTSVTNVGPRVGLGLP  
 LSPRLLPTATTNGHPQLPGHAKPGTPA  
 LETLETTTPAMAAPKDDGFLNGSCSGLDEEASGPERPPALLQEEWETVM

As mentioned above, further exemplifying PTH/PTHrP receptor chameleons have been constructed. These constructs, in particular relate to amino acid sequence junction between PTH/PTHrP receptor's carboxy terminal tail and YFP for various PTH receptor-cam constructs.

The best FRET efficiency are in the following order: PTHR-cam8>PTHR-cam9>PTHR-cam7>PTHR-cam2>PTHR-cam5>PTHR-cam1>PTHR-cam4>PTHR-cam3=PTHR-cam10. PTH mediated a FRET decrease in these constructions.

PTHR-cam8 is the construct described above in SEQ ID NOS: 13 and 14.

Further constructs are defined below. These constructs correspond to PTHR-cam8, yet, they comprise a modified C-terminus.

**SEQ ID NO: 29: PTHR-cam7 amino acid sequence:**

EVQAEIKKSWSRWTLALDFKRKARSMVSKGEELFTGVVPILVELDGDVNGHKFS  
VSGEGEGDATYGKLTCLKFICTTGKLPVPWPTLVTT  
FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY  
NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALSKDPNEKRDHMLLEF  
VTAAGITLGMDELYK

**SEQ ID NO: 30: PTHR-cam9 amino acid sequence:**

EVQAEIKKSWSRWTLALDFKRKARSGSSSYSMVSKGEELFTGVVPILVELDGDV  
NGHKFSVSGEGEGDATYGKLTCLKFICTTGKLPVPWPTLVTT  
FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY  
NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALSKDPNEKRDHMLLEF  
VTAAGITLGMDELYK

**SEQ ID NO: 31: PTHR-cam8 amino acid sequence:**

EVQAEIKKSWSRWTLALDFKRKARSGSSSYSYGMVSKGEELFTGVVPILVELDG  
DVNGHKFSVSGEGEGDATYGKLTCLKFICTTGKLPVPWPTLVTT  
FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY  
NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALSKDPNEKRDHMLLEF

VTAAGITLGMDELYK

SEQ ID NO: 32: PTHR-cam2 amino acid sequence:

EVQAEIKKSWSRWTLALDFKRKARSGSSSYSGPMVMVSKGEELFTGVVPILVE  
LDGDVNGHKFSVSGEGEGDATYGKLTTKFICTTGKLPVPWPTLVTT  
FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY  
NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALSKDPNEKRDHMLLEF  
VTAAGITLGMDELYK

SEQ ID NO: 33: PTHR-cam5 amino acid sequence:

EVQAEIKKSWSRWTLALDFKRKARSGSSSYSGPMVSHTSVTNVGPRVGLMVS  
KGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTTKFICTTGKLPVP  
WPTLVTT  
FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY  
NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALSKDPNEKRDHMLLEF  
VTAAGITLGMDELYK

SEQ ID NO: 34: PTHR-cam1 amino acid sequence:

EVQAEIKKSWSRWTLALDFKRKARSGSSSYSGPMVSHTSVTNVGPRVGLGLP  
 LSPRLLPTATTNGHPQLPGHAKPGTPA  
 LETLETMVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTTKFI  
CTTGKLPVPWPTLVTT  
FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY

NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALS KDPNEKRDH MVLLEF  
VTAAGITLGMD ELYK

SEQ ID NO: 35: PTHR-cam4 amino acid sequence:

EVQAEIKKSWSRWTLALDFKRKARSGSSSYSGPMVSHTSVTNVGPRVGLGLP  
 LSPRLLPTATTNGHPQLPGHAKPGTPA  
 LETLETPPAMAAPKDDGFL MVSKGEELFTGVVPILVELDGDVNGHKFSVSGEG  
EGDATYGKLT LKFICTTGKLPVPWPTLVTT  
FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY  
NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALS KDPNEKRDH MVLLEF  
VTAAGITLGMD ELYK

SEQ ID NO: 36: PTHR-cam3 amino acid sequence:

EVQAEIKKSWSRWTLALDFKRKARSGSSSYSGPMVSHTSVTNVGPRVGLGLP  
 LSPRLLPTATTNGHPQLPGHAKPGTPA  
 LETLETPPAMAAPKDDGFLNGSCSGLDEEASGPE MVSKGEELFTGVVPILVEL  
DGDVNGHKFSVSGEGEGDATYGKLT LKFICTTGKLPVPWPTLVTT  
FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY  
NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALS KDPNEKRDH MVLLEF  
VTAAGITLGMD ELYK

SEQ ID NO: 37: PTHR-cam10 amino acid sequence :

EVQAEIKKSWSRWTLALDFKRKARSGSSSYSGPMVSHTSVTNVGPRVGLGLP  
 LSPRLLPTATTNGHPQLPGHAKPGTPA

LETLETPPAMAAPKDDGFLNGSCSGLDEEASGPERPPALLQEEWETVMMVSK  
GEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTGKLPVPW  
PTLVTT  
FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY  
NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALS KDPNEKRDHMLLEF  
VTAAGITLGMDLYK

The following control construct is exemplified: Amino acid sequence junction between PTH/PTHrP receptor's carboxy terminal tail and YFP and CFP in PTHR-FRET<sup>control</sup>.

This construction gave a strong FRET efficiency. However, PTH, as ligand, does not mediated a change in FRET in this construction.

SEQ ID NO: 38: PTHR- PTHR-FRET<sup>control</sup> amino acid sequence:

EVQAEIKKSWSRWTLALDFKRKARMVSKGEELFTGVVPILVELDGDVNGHKFSV  
SGEGEGDATYGKLTCLKFICTTGKLPVPWPTLVTT  
FGYGLQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFE  
GDTLVNRIELKGIDFKEDGNILGHKLEY  
NYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPD  
NHYLSYQSALS KDPNEKRDHMLLEF  
VTAAGITLGMDLYKSGSSSYSGPMVSHTSMVSKG  
EELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTGKLPVPWP  
TLVTTLTWGVQCFSRYPDHMKQHDF  
KSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILG  
HKLEYNYISHNVYITADKQKNGIKAN  
FKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSALS KDPNEKRDH  
MVLEFVTAAGITLGMDLYK

### Example II: Generation of receptor constructs with agonist-sensitive FRET

A series of receptor constructs were generated that carried CFP and/or YFP in various positions of the third intracellular loop and/or of the carboxy-terminus of the PTHR and the  $\alpha_{2A}$ AR, respectively. The two constructs showing the best cell surface expression, pharmacological properties and agonist sensitivity were further used in this study and are referred to as receptor chameleons PTHR-cam and  $\alpha_{2A}$ AR-cam (Fig. 1A).

Emission fluorescence spectra were recorded from HEK293 cells stably expressing "PTHR-cam8" and various control constructs. The sequences representing "PTHR-cam8" are represented in appended SEQ ID NOS: 13 and 14. Excitation of a PTHR carrying only a CFP-moiety in its 3<sup>rd</sup> intracellular loop (PTHR-CFP<sub>3-loop</sub>) with light at 436 nm resulted in an emission at 480 nm (corresponding to the CFP emission, Fig. 1B). The additional presence of YFP in the carboxy terminus (PTHR-cam) lead to a reduced emission at 480 nm plus a strong emission at the characteristic wavelength of YFP (535 nm, Fig. 1B). Photobleaching of the acceptor confirmed that the latter emission was primarily due to FRET (see Fig. 1C). The presence of YFP in the C-terminus alone (PTHR-YFP<sub>C-term</sub>) did not result in significant emission at 535 nm when excited at 436 nm (Fig. 1B). Receptor constructs containing only YFP showed no specific emission peak at 535 nm when excited at 436 nm (Fig. 1B). It is of note that further PTHR-cam constructs have been prepared which are shown in related SEQ ID NOS: 29 to 37. A control construct is shown in SEQ ID NO: 38.

Signals recorded from single HEK293 cells expressing PTHR-cam8 were then analysed at emissions of 480 nm (CFP) and 535 nm (YFP) upon excitation at 436 nm (CFP excitation). The microscopic illumination allowed photobleaching experiments in order to verify that the emission at 535 nm was indeed due to FRET. After bleaching of the acceptor in the PTHR-cam8 construct with intense light at 480 nm, the emission at 480 nm increased by 50±3% together with a more than 5-fold reduction of the 535 nm emission (Fig. 1C). Similar spectral and

photobleaching data were obtained with the " $\alpha_{2A}$ AR-cam" stably expressed in HEK293 cells. The corresponding constructs for " $\alpha_{2A}$ AR-cam" are depicted in SEQ ID NOS: 11 and 12.

The effects of the agonist PTH on the FRET signal of PTHR-cam were investigated, measured as the background-corrected emission intensity ratio  $F_{535}^*/F_{480}^*$ . After addition of 1  $\mu$ M PTH, the ratio  $F_{535}^*/F_{480}^*$  rapidly decreased (Fig. 1D). After a short delay ( $\approx 600$  ms) the decrease followed a mono-exponential time-course with a time-constant  $\tau = 3.00 \pm 0.25$  s ( $n=9$ ). The symmetrical increase in CFP emission and decrease in YFP emission indicate that the change was due to a decrease in FRET. Control experiments with co-expression of PTHR-CFP<sub>3-loop</sub> and PTHR-YFP<sub>C-term</sub> showed no FRET in the absence or in the presence of PTH (1  $\mu$ M) and thus made it unlikely that the signals resulted from *intermolecular* FRET in receptor dimers. It should be noted, however, that the *intramolecular* nature of this signal does not exclude the presence of receptor dimers.

Similarly,  $\alpha_{2A}$ AR-cam showed intramolecular FRET, and again the specific agonist, i.e. noradrenaline, caused a decline of the FRET signal (see Figs. 3, 4, 6). Virtually identical results were obtained with the two receptors expressed in other cell lines (CHO, PC12; data not shown). The agonist-induced decreases in FRET in the two types of receptors suggest that the agonist-induced conformational switch is similar in class 1 and class 2 GPCRs. Because of the location of the CFP and YFP in the receptors (Fig. 1A) they are compatible with a movement of the 3<sup>rd</sup> intracellular loop away from the C-terminus as predicted by computer simulations of the  $\alpha_{1B}$ -adrenergic receptor (Greasley, J. Biol. Chem. 276 (2001), 46485-46494).

### **Example III: Pharmacological characterization of the receptor-chameleon**

The receptor-chameleon constructs stably expressed in HEK293 cells retained the typical ligand binding which was of somewhat lower affinity than for the corresponding wild-type receptors (Fig. 2): the PTH affinity was  $K_i = 15.5 \pm 0.9$  nM

for PTHR-cam and  $K_i=2.4\pm0.4$  nM for PTHR; the nordrenaline affinity was  $K_i=5.0\pm0.8$  for  $\alpha_{2A}AR$  and  $K_i=16.7\pm1.4$   $\mu M$  for  $\alpha_{2A}AR$ -cam. Note that the insertion of the GFP variants into the third loop and carboxy terminus of the receptor might cause a conformational destabilization of the receptor resulting in a deviation in the binding properties of the GPCRs-cam. Such destabilization might result in a faster activation switch of the receptor. However, the  $\alpha_{2A}AR$ -cam induced GIRK current activation was not faster than the wild type  $\alpha_{2A}AR$  (Bünemann, (2001), loc. cit.).

PTHR-cam and  $\alpha_{2A}AR$ -cam signalled efficiently to adenylyl cyclase ( $EC_{50}=12.8\pm1.4$  nM) and to the GIRK channel ( $EC_{50}=1.08\pm0.01$   $\mu M$ ), respectively (Fig. 2A-B). A difference in the signal amplification between PTHR wild type- and PTHR-cam receptors (1  $\mu M$  PTH mediated 28 vs. 11.5-fold increase in cAMP in HEK-293 cells expressing PTHR or PTHR-cam, respectively) is in part due to the higher expression level of the wild type receptors ( $1.01\times10^6$  vs  $0.34\times10^6$  receptors/cells for PTHR and PTHR-cam, respectively). Finally, confocal microscopy showed that the majority of PTHR-cam and  $\alpha_{2A}AR$ -cam receptors in stably transfected cells were correctly present at the cell surface (Fig. 2C). These data indicate that both receptor constructs were properly targeted to the cell surface upon expression in HEK293 cells and retained essential binding properties as well as significant G-protein-mediated signalling of the corresponding wild-type receptors. Similar data were obtained upon expression in various cell lines (e.g. CHO or PC12).

#### **Example IV: The agonist-mediated FRET signal is coupled to receptor activation**

It was verified that the agonist-induced changes in FRET does indeed reflect the conformational change of the receptor. The following verification experiments were carried out.



First, it was shown that the FRET signal was indeed caused by the receptors themselves and not by interactions with other proteins such as G-proteins or  $\beta$ -arrestins. Therefore, we studied PTHR-cam under conditions that exclude interactions with these proteins. In isolated cell membranes prepared from HEK293 cells stably expressing PTHR-cam – i.e. in the absence of cytosolic proteins – the PTH-induced signal had the same magnitude as in intact cells (Fig. 3A). Further stripping the cell membranes with 6 M urea – a treatment known to leave GPCRs intact but to denature virtually all other proteins (Sheikh, J. Biol. Chem. 274 (1999), 17033-17041; Lim, Biochem. J. 354 (2001), 337-344) – did also not affect the magnitude of the PTH-induced signal at saturating concentrations (Fig. 3A). And finally, inactivating  $G_i$  and  $G_o$  with pertussis toxin in cells expressing the  $\alpha_{2A}$ AR-cam did not affect the noradrenaline-induced FRET signal, indicating that the signal was not due to a receptor/G-protein interaction. Taken together, these data document that the FRET signals were not caused by interactions of the receptors with other proteins.

Second, it was shown that the FRET signals correspond to the activation state of the receptor. Accordingly, the effects of agonists and antagonists were tested. A truncated variant of PTH, PTH7-34, which is a low affinity antagonist, failed to induce a change in FRET (Fig. 3A). Similarly, noradrenaline (10  $\mu$ M) induced a rapid decrease of the FRET signal in the  $\alpha_{2A}$ AR-cam (Fig. 3B), while saturating concentrations of the high affinity  $\alpha_2$ -adrenergic receptor antagonist phentolamine (10  $\mu$ M) did not alter the FRET signal when given alone. However, phentolamine rapidly reverted the noradrenaline-induced signal (Fig. 3B). This is compatible with its nature as a competitive antagonist. Thus, the rigorous agonist dependence on the change of the FRET signal mirrors the active state of the receptor.

Third, binding of G-proteins to receptors is known to enhance formation of the active, agonist-bound state. Because of the reduced ability of the receptor chameleons to couple to G-proteins, such assays required the addition of exogenous G-proteins. After addition of purified  $G_o$  to membranes containing  $\alpha_{2A}$ AR-cam, the agonist [ $^3$ H]UK14304 bound with high affinity to the receptors

( $K_d=3.4\pm0.8$  nM; Supplementary Fig. 1). Addition of the stable GTP-analog GTP $\gamma$ S reduced this affinity ( $K_d=9.6\pm1.1$  nM) indicative of a disruption of the high-affinity receptor/G-protein complex. GTP $\gamma$ S reduced the binding of 5 nM [ $^3$ H]UK14304 by more than 50% (Fig. 3C *right* panel). Similarly, GTP $\gamma$ S reduced the FRET signals caused by a 5 nM UK14304 in the same membrane preparation (i.e. in the presence of  $G_o$ ) by more than 50% (Fig. 3C *left* panel). GTP $\gamma$ S did not affect the signal of saturating concentrations of UK14304, which is in agreement with the lack of effect of urea-treatment on the maximal PTH-induced signal obtained in membranes (Fig. 3A). Taken together, these data suggest that the FRET-signal originates in the active conformation of the receptor itself, and that this active conformation binds to and is stabilized by G-proteins.

#### **Example V: FRET changes mediated by a partial agonist**

The FRET assay also properly reflected partial agonism (Fig. 4): Compared to the full agonist noradrenaline, the high affinity partial agonist clonidine at saturating concentrations (10  $\mu$ M) gave a three-fold smaller FRET signal (Fig. 4). Subsequent application of noradrenaline (10  $\mu$ M) still produced the full response. The simultaneous addition of clonidine (10  $\mu$ M) restored this response back to the partial response seen with clonidine alone; and again, after washout, noradrenaline still produced the full initial response. These data correspond exactly to the predicted properties of a high affinity partial agonist. However, compared to other assays used so far to detect partial agonism, the FRET assay is not dependent on transducer and effector proteins. Instead it reflects directly the partial agonist effects on the receptors themselves. Mechanistically, the ability of clonidine to partially reverse the agonist-mediated signal suggests that the partial agonist restrains the complete movement between the 3<sup>rd</sup> intracellular loop and the C-terminus. This is compatible with the notion that the partial agonism process occurs at the receptor level by inducing a restrained conformational change within the agonist binding site (Ghanouni, J. Biol. Chem. 276 (2001), 24433-24436).

### Example VI: Comparing receptor activation with desensitization

Receptor activation should precede receptor deactivation. Therefore, we measured the rate-limiting step in PTHR-deactivation (Villardaga, (2002), loc. cit.; Castro, *Endocrinology* 143 (2002), 3854-3865), the association of  $\beta$ -arrestin with the receptor, again with a FRET-based approach. To this end, we co-expressed functional PTHR carrying CFP at its C-terminus (PTHR-CFP<sub>C-term</sub>) and  $\beta$ -arrestin2 fused at its C-terminus to YFP ( $\beta$ -arrestin2-YFP). Measuring the appearance of FRET between the CFP and the YFP then monitored PTH-induced binding of  $\beta$ -arrestin2 to the receptor. The dynamics of this signal were compared with the receptor activation of PTHR-cam (Fig. 5). The initial ratio  $F_{535}^*/F_{480}^*$  was  $1.10 \pm 0.05$  for cells co-expressing PTHR-CFP<sub>C-term</sub> and  $\beta$ -arrestin2-YFP. After addition of PTH (100 nM), the ratio increased by up to  $32 \pm 11$  % with a  $t_{1/2}$  of  $150 \pm 12.1$  s ( $n=8$ ), reflecting the PTH-mediated receptor/ $\beta$ -arrestin2 association. The same concentration of PTH (100 nM) had a 5-fold faster effect on PTHR-cam, with a  $t_{1/2}$  of  $32 \pm 1.9$  s ( $n=4$  experiments). Furthermore, the lag time between addition of PTH and the beginning of the response was about three times shorter for the activation signal (PTHR-cam) than for  $\beta$ -arrestin translocation (Fig. 5, Inset). Thus, the signal for receptor activation does indeed begin earlier and proceeds much faster than that for receptor deactivation.

### Example VII: Differential speed of activation between hormone and neurotransmitter receptors

Time-resolved determination of the FRET signals recorded from single cells after activation with various concentrations of PTH and noradrenaline, respectively, allowed the analysis of the switch kinetics (Fig. 6). Under all conditions, the decrease of the ratio  $F_{535}^*/F_{480}^*$  followed a monoexponential time-course. Increasing concentrations of agonist resulted in shorter delay times as well as faster time-courses of the signals. At low agonist concentrations, the rate constants ( $k_{obs}$ ) increased in proportion to agonist concentration (Fig. 6B), indicating that agonist binding to the receptors was the rate-limiting step. At higher

concentrations of agonist, the rate constants reached a maximum, suggesting that a step other than the collisional probability of agonist/receptor became rate-limiting. This limit is not due to technical limitations of the system, which allows complete solution exchange in less than 10 ms. Thus, it is most likely the agonist-mediated conformational switch of the receptors. The time constant required for the receptor activation was less than 40 ms in the case of the  $\alpha_{2A}$ AR-cam. This is more than 5000 times faster than the activation time measured for chemically labelled, purified  $\beta_2$ -adrenergic receptors (Gether, *J. Biol. Chem.* 270 (1995), 28268-28275; Jensen, *J. Biol. Chem.* 276 (2001), 9279-9290; Ghanouni, (2001), loc. cit.; Ghanouni, *Proc. Natl. Acad. Sci. USA* 98 (2001), 5997-6002) and corresponds well to the physiological requirements of neurotransmitter receptors. In contrast, the kinetics of the FRET-signal were 25-fold slower ( $\tau \approx 1$ s) for the PTH-receptor. This may be explained by the fact that PTH is a large agonist and that its binding appears to involve several contact points both in the extended N-terminus and in the core of the PTHR (Gardella, *Trends Endocrinol. Metabolism* 12 (2001), 210-217).

The saturation of the  $k_{obs}$ -values together with the concentration-dependent delay times (Fig. 6) are compatible with a simple two-step process of first-order agonist binding and subsequent receptor activation. The delay times indicate that the FRET signal reflects the conformational switch underlying receptor activation and not just receptor-agonist contact. Mechanistically, this switch appeared to be similar in a class 1 and a class 2 GPCR, but the activation was much faster in the class 1  $\alpha_{2A}$ AR-cam. The slower activation of the class 2 PTH receptor is compatible with the slow hormonal effects of PTH, compared with the fast synaptic action of noradrenaline. Millisecond switch times have so far been thought to be limited to ion channel receptors (Chang, *Nature Neurosci.* 5 (2002), 1163-1168) or to rhodopsin (Okada, *Trends Biochem. Sci.* 26 (2001), 318-324). However, data presented herein document that also a "neurotransmitter GPCR" can be switched in the millisecond time scale, and it is shown that the extent said switching is dependent on the intrinsic efficacy of the ligand.

**Example VIII: A2A adenosine receptor activation determined by intramolecular FRET:**

Recombinant YFP and eCFP were fused to recombinant human A2A adenosine receptors, similar as described for  $\alpha$ 2A adrenergic receptors and PTH/PTHrP receptors using conventional cloning strategies. Specifically amino acids between amino acid 14 after the 5<sup>th</sup> transmembrane helix and amino acid 10 proximal to the 6<sup>th</sup> transmembrane helix were replaced by eCFP (corresponding to the third intracellular loop). The c-terminus of the receptor was truncated after amino acid 33 and was fused to YFP.

HEK293 cells were transiently transfected with A2A-CFP14/10-YFPC33 (exemplified constructs shown in SEQ ID NOS: 15 and 16) using effectene (Qiagen) according to the manufacturers protocol. 12h post transfection cells were replated on glass coverslips coated with poly-L-lysine and cultured for another 12-36 h. Glass coverslips were subsequently washed with HEPES buffered solution and placed under an inverted fluorescence microscope (Zeiss axiovert 135). Using excitation light of 436 nm (bandwidth 10 nm) cells expressing fluorescent receptors were excited and emission of single cells was detected at 480 nm ( $480 \pm 20$  nm) for CFP-emission and 535 nm ( $535 \pm 15$  nm) using a dual wavelength detection system equipped with amplified photodiodes (Till photonics). CFP and YFP emission in response to 10 ms illumination periods every 100 ms was recorded and after correcting for bleed through of the CFP emission into the 535 nm channel a ratio of YFP/CFP emission was calculated (nFRET).

Cells were continuously superfused with control solution (in mM : 140 NaCl; 5.4 KCl; 2 CaCl<sub>2</sub>; 1 mM MgCl<sub>2</sub>; 10 mM Glucose and 10 HEPES/NaOH pH 7.3) using a fast solenoid valve operated superfusion system ALA-VM8 (ALA Scientific Instruments). Upon superfusion with agonist containing solution (100  $\mu$ M adenosine) a rapid 1.5-2% decrease in nFRET was detected, less in amplitude, but otherwise similarly as described for  $\alpha$ 2A-cam and PTHR-cam8. After withdrawal of agonist, nFRET increased to the initial value within 5-10 s.

These results indicate that agonist induced conformational changes of A2A adenosine receptors can be detected by changes in intramolecular FRET of A2A-CFP14/10-YFPC33 receptors, similar as described for  $\alpha$ 2A-cam and PTHR-cam8.

**Example IX: Inactivation of human G protein $\alpha$  ( $G\alpha$ ) by introduction of a fluorophore**

Human  $G\alpha$ s was mutated by introducing of the YFP into the loop  $\alpha$ A- $\alpha$ B at the position 116 and was coexpressed with  $G\beta\gamma$  subunits of heterotrimeric G proteins. Using fluorescence microscopy no membrane-targeting of  $G\alpha$ s was observed. Instead the fluorescent protein was found in perinuclear spots, most likely resembling lysosomes. The introduction of the YFP tag into  $G\alpha$ s led to a severe defect of either protein folding or targeting of this fusion protein in intact cells. In addition no FRET between fluorescent  $G\beta\gamma$ -CFP subunits and  $G\alpha$ -YFP was detected, indicating no proper interaction of these natural binding partners.

**Claims**

1. Recombinant seven-transmembrane receptor, whereby the amino terminus of said recombinant receptor is located on an extracellular side and the carboxy-terminus is located on an intracellular side of a membrane, comprising at least two detectable labels, whereby a first of said at least two detectable labels is or is located (on) the carboxy-terminus and whereby a second of said at least two labels is or is located (on) the first or third intracellular loop or whereby a first of said at least two labels is or is located (on) the first intracellular loop and a second of at said at least two labels is or is located (on) the third intracellular loop.
2. Recombinant membrane receptor of claim 1, whereby said first label is or is located (on) the third intracellular loop of said membrane receptor and wherein said second label is or is located (on) the carboxy terminus.
3. Recombinant membrane receptor of claim 1 or 2, whereby said membrane receptor is a G-protein-coupled receptor or a proto-oncogene.
4. Recombinant membrane receptor of claim 1, whereby the proto-oncogene is Smoothed receptor (Smo) or whereby said G-protein-coupled receptor (GPCR) is selected from the group consisting of a rhodopsin/ $\beta$ 2 adrenergic receptor-like GPCR, a glucagon/VIP/calcitonin receptor-like GPCR and a metabotropic neurotransmitter/calcium receptor.
5. Recombinant membrane receptor of claim 4, whereby said rhodopsin/ $\beta$ 2-adrenergic receptor-like GPCR is the  $\alpha$ 2A adrenergic receptor or the adenosine receptor A2A or wherein said glucagon/VIP/calcitonin receptor-like GPCR is the parathyroid hormone (PTH) receptor.

6. Recombinant membrane receptor of any one of claims 3 to 5, whereby said G-protein-coupled receptor or said proto-oncogene is of human or of mouse origin.
7. Recombinant membrane receptor of any one of claims 1 to 6, whereby said detectable labels are fluorescent labels or bioluminescent labels.
8. Recombinant membrane receptor of claim 7, whereby said fluorescence labels are selected from the group consisting of GFP, YFP, CFP, BFP, citrine, sapphire and dsRed, whereby said bioluminescent labels is luciferase (like renilla luciferase or firefly luciferase), or whereby said fluorescence label is produced by binding the Flash compound to specific epitopes of said 1<sup>st</sup> and 3<sup>rd</sup> loop or said C-terminus of the recombinant seven-transmembrane receptor.
9. Recombinant membrane receptor of any one of claims 3 to 8, whereby said G-protein-coupled receptor comprising at least two labels is selected from the group consisting of:
  - (a) a polypeptide as shown in SEQ ID NOS: 12, 14 or 16;
  - (b) a polypeptide encoded by a nucleic acid sequence as depicted in any one of SEQ ID NOS: 11, 13 or 15;
  - (c) a recombinant membrane receptor of any one of claims 3 to 8 encoded by a nucleotide sequence which hybridizes to a nucleotide sequence as defined (b); and
  - (d) a recombinant membrane receptor of any one of claims 3 to 8 encoded by a nucleic acid sequence being degenerate as a result of the genetic code to a nucleic acid sequence as defined in (b) or (c).
10. Recombinant membrane receptor of any one of claims 3 to 9, wherein the third intracellular loop being or comprising said first label is selected from the group consisting of
  - (a) a polypeptide depicted in SEQ ID NOS: 18, 22 or 26;



- (b) a polypeptide encoded by a nucleic acid sequence as depicted in SEQ ID NOS: 17, 21 or 25;
  - (c) a third intracellular loop encoded by a nucleotide sequence which hybridizes to a nucleotide sequence as defined (b); and
  - (d) a third intracellular loop encoded by a nucleic acid sequence being degenerate as a result of the genetic code to a nucleic acid sequence as defined in (b) or (c).
- 11. A nucleic acid molecule encoding the recombinant seven-transmembrane receptor of any one of claims 1 to 10.
  - 12. A vector comprising a nucleic acid molecule as defined in any one of claims 8 to 11.
  - 13. The vector of claim 12, which is an expression vector.
  - 14. A host transformed with a vector of claim 12 or 13 or transfected with a nucleic acid molecule as defined in any one of claims 8 to 11.
  - 15. The host of claim 14 which is a mammalian cell, an amphibian cell, a fish cell, an insect cell, a fungal cell, a plant cell or a bacterial cell, or a transgenic non-human animal.
  - 16. The host of claim 15, wherein said mammalian cell is a CHO-cell or a HEK293 cell, a PC12 cell, a (primary) cardiomyocyte or a cultured neuronal cell.
  - 17. The host of claim 15, wherein said amphibian cell is an oocyte, preferably a xenopus oocyte.

18. A method for producing a recombinant membrane receptor as defined in any one of claims 1 to 10 comprising culturing/raising the host of any one of claims 14 to 17 and, optionally, isolating the produced polypeptide.
19. A method for identifying molecules or compounds which are capable of activating, deactivating or inactivating the (biological/pharmacological) function of a seven-transmembrane receptor, comprising the steps of
  - (a) contacting the recombinant membrane receptor as defined in any one of claims 1 to 10 or a host or a host cell of any one of claims 14 to 17 with (a) molecule(s) or compound(s) to be tested; and
  - (b) measuring whether said molecule(s) or compound(s) to be tested lead(s) to a modification of a signal provided by said at least two detectable labels.
20. A method of screening for molecules or compounds which are activators (agonists) or inhibitors (antagonists) of the (biological/pharmacological) function of a seven-transmembrane receptor comprising the steps of
  - (a) contacting a recombinant membrane receptor as defined in any one of claims 1 to 10 or a host or a host cell of any one of claims 14 to 17 with the molecule or compound to be tested;
  - (b) measuring and/or detecting a response comprising a modification of a signal provided by said at least two detectable labels; and
  - (c1) comparing said response to a standard response as measured in the absence of said candidate molecule/compound
  - (c2) comparing said response to the response of a control membrane receptor which comprises at least two detectable labels on the C-terminus; or
  - (c3) comparing said response to control seven-transmembrane receptor which comprises only one detectable label.

21. A method for identifying molecules or compounds which are capable of eliciting a (biological/pharmacological) response of a seven-transmembrane protein, comprising the steps of
  - (a) contacting a membrane protein of any one of claims 1 to 10, or a host or host cell of claims 14 to 17 with the molecule or compound to be tested; and
  - (b) identifying among these molecules/compounds the molecules/compounds which are capable of eliciting a change in energy emitted by said at least two detectable labels comprised on the recombinant membrane receptor as defined in any one of claims 1 to 10.
22. The method of any one of claims 19 to 21, whereby said response or said energy change is an increase or a decrease of fluorescence resonance energy transfer (FRET).
23. The method of any one of claims 19 to 21, whereby said response or said energy change is an increase or a decrease of bioluminescent resonance energy transfer (BRET).
24. A diagnostic composition comprising the recombinant membrane protein of any one of claims 1 to 10 or the nucleic acid molecule as defined in claims 8 to 11, the vector of claims 12 or 13, the host cell of claims 14 to 17 or organs or cells of the non-human transgenic animal of claim 15.
25. A kit comprising the recombinant membrane protein of any one of claims 1 to 10 or the nucleic acid molecule as defined in claims 8 to 11, the vector of claims 12 or 13, the host cell of claims 14 to 17 or organs or cells of the non-human transgenic animal of claim 15.
26. Use of the recombinant membrane protein of any one of claims 1 to 10 or the nucleic acid molecule as defined in claims 9 or 10, the vector of claims

11 or 12, the host cell of claims 13 to 16 or organs or cells of the non-human transgenic animal of claim 16 for the detection of (a) modifier(s) of the biological activity of seven-transmembrane receptors in vivo or in vitro.

**Abstract**

The present invention relates to recombinant seven-transmembrane receptor, whereby the amino terminus of said recombinant receptor is located on an extracellular side and the carboxy-terminus is located on an intracellular side of a membrane, comprising at least two detectable labels, whereby a first of said at least two detectable labels is or is located (on) the carboxy-terminus and whereby a second of said at least two labels is or is located (on) the first or third intracellular loop or whereby a first of said at least two labels is or is located (on) the first intracellular loop and a second of at said at least two labels is or is located (on) the third intracellular loop. Furthermore, nucleic acid molecules encoding said recombinant seven-transmembrane receptors are described as well as vector and host cells comprising the same. Furthermore, the present invention provides for identification and screening methods for molecules or compounds which are capable of modifying the biological end of pharmacological function of seven-transmembrane receptor proteins. Finally, diagnostic compositions comprising the compounds of the present invention as well as kits comprising said compounds are disclosed.

Figure 1




Figure 1 part 2




Figure 2

3/7






Figure 3




Figure 4




Figure 5





Figure 6



PCT Application  
**PCT/EP2003/014679**



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**